

## Annex

## **KEY CONDOM USE INDICATORS**

Table displays most recent figure per indicator from relevant source.	Botswana	Burkina Faso	Cameroon	eSwatini	Kenya	Malawi	Mozambique	Nigeria	South Africa	Tanzania	Uganda	Zambia	Zimbabwe
% Condom use at last NMNC sex (Women 15-49)	<b>82</b> % (2013)	<b>59%</b> (2010)	<b>55%</b> (2011)	<b>54%</b> (2007)	<b>57%</b> (2014)	<b>50%</b> (2015)	<b>42</b> % (2015)	<b>40%</b> (2013)	<b>58%</b> (2016)	<b>51</b> % (2012)	<b>38</b> % (2016)	<b>41</b> % (2013)	<b>67%</b> (2015)
% Condom use at last NMNC sex (Men 15-49)	<b>82%</b> (2013)	<b>74%</b> (2010)	<b>74%</b> (2011)	<b>67%</b> (2007)	<b>76</b> % (2014)	<b>76%</b> (2015)	<b>47%</b> (2015)	<b>58%</b> (2013)	<b>65%</b> (2016)	<b>60%</b> (2012)	<b>41%</b> (2016)	<b>56%</b> (2013)	<b>85</b> % (2015)
% Condom use during premarital sex (Women 15-24)		<b>62</b> % (2010)	<b>59%</b> (2011)	<b>54</b> % (2007)	<b>61</b> % (2014)	<b>54%</b> (2015)	<b>56%</b> (2015)	<b>44%</b> (2013)		<b>37%</b> (2016)	<b>45</b> % (2016)	<b>40</b> % (2014)	<b>50%</b> (2015)
% Condom use during premarital sex (Men 15-24)		<b>76</b> % (2010)	<b>72</b> % (2011)	<b>70</b> % (2007)	<b>75</b> % (2014)	<b>73</b> % (2015)	<b>47</b> % (2015)	<b>58%</b> (2013)		<b>41</b> % (2016)	<b>58%</b> (2016)	<b>49</b> % (2014)	<b>81%</b> (2015)
% Condom use at last paid sexual intercourse (Men)		<b>33</b> % (2010)	<b>81</b> % (2011)		<b>74</b> % (2014)	<b>74</b> % (2015)	<b>31%</b> (2015)	<b>66%</b> (2013)		<b>59</b> % (2010)	<b>73</b> % (2016)	<b>60%</b> (2014)	<b>90%</b> (2015)
% Condom use at last anal sex (MSM)	<b>67%</b> (2013)		<b>78%</b> (2018)	<b>67%</b> (2018)	<b>80%</b> (2018)	<b>63</b> % (2018)	<b>31%</b> (2018)	<b>83%</b> (2018)	<b>81</b> % (2018)	<b>14%</b> (2018)	<b>39%</b> (2018)		
HIV prevalence	<b>19</b> % (2013)	<b>1</b> % (2016)	<b>4</b> % (2011)	<b>27</b> % (2017)	<b>6</b> % (2008)	<b>9</b> % (2015)	<b>13</b> % (2015)	<b>3%</b> (2017)	<b>19%</b> (2017)	<b>5</b> % (2011)	<b>7</b> % (2011)	<b>13</b> % (2013)	<b>14%</b> (2015)
% w knowledge of a formal source for condoms (Women 15-24)		<b>76%</b> (2010)	<b>71%</b> (2011)	<b>85</b> % (2007)	<b>71%</b> (2014)	<b>79%</b> (2010)	<b>65%</b> (2011)	<b>46</b> % (2013)		<b>65%</b> (2012)	<b>75%</b> (2011)	<b>84%</b> (2013)	<b>48</b> % (2015)
% w knowledge of a formal source for condoms (Men 15-24)		<b>90%</b> (2010)	<b>87</b> % (2011)	<b>88</b> % (2007)	<b>88</b> % (2014)	<b>89%</b> (2010)	<b>91%</b> (2011)	<b>68%</b> (2013)		<b>85</b> % (2012)	<b>90%</b> (2011)	<b>94%</b> (2013)	<b>86</b> % (2015)
% w comprehensive AIDS knowledge (Women 15-24)	<b>48</b> % (2013)	<b>31%</b> (2010)	<b>27</b> % (2011)	<b>52</b> % (2007)	<b>57</b> % (2014)	<b>42</b> % (2010)	<b>31%</b> (2015)	<b>24</b> % (2013)		<b>40</b> % (2012)	<b>39%</b> (2016)	<b>42</b> % (2014)	<b>46</b> % (2015)
% w comprehensive AIDS knowledge (Men 15-24)	<b>48</b> % (2013)	<b>36%</b> (2010)	<b>34</b> % (2011)	<b>52%</b> (2007)	<b>64</b> % (2014)	<b>45</b> % (2010)	<b>30%</b> (2015)	<b>34</b> % (2013)		<b>47%</b> (2012)	<b>40</b> % (2011)	<b>47%</b> (2014)	<b>47%</b> (2015)
Total condoms dist'd (M)	<b>28</b> (2016)	<b>0.5</b> (2017)	<b>60</b> (2016)	<b>19</b> (2017)	<b>234</b> (2017)	<b>32</b> (2017)	<b>186</b> (2016)	<b>363</b> (2016)	<b>1,008</b> (2017)	<b>140</b> (2016)	<b>206</b> (2016)	<b>36</b> (2015)	<b>123</b> (2017)
Free condoms dist'd (M)	<b>23</b> (2016)		<b>16</b> (2016)	<b>18</b> (2017)	<b>199</b> (2017)	<b>14</b> (2017)	<b>58</b> (2016)	<b>115</b> (2016)	<b>917</b> (2017)	<b>38</b> (2016)	<b>183</b> (2016)	<b>25</b> (2015)	<b>105</b> (2017)
SM sector condoms dist'd (M)	<b>0.0</b> (2016)	<b>0.5</b> (2017)	<b>20</b> (2016)	<b>1.1</b> (2017)	<b>33</b> (2017)	<b>18</b> (2017)	<b>16</b> (2016)	<b>174</b> (2016)	<b>0.0</b> (2017)	<b>99</b> (2016)	<b>22</b> (2016)	<b>6.2</b> (2015)	<b>16</b> (2017)
Commercial sector condoms dist'd (M)	<b>5.1</b> (2016)		<b>24</b> (2016)		<b>2.7</b> (2017)		<b>17</b> (2016)	<b>75</b> (2016)	<b>91</b> (2017)	<b>2.9</b> (2016)	<b>1.0</b> (2016)	<b>5.0</b> (2015)	<b>1.5</b> (2017)
Condoms per male	<b>41</b> (2016)	<b>0.1</b> (2017)	<b>9.5</b> (2016)	<b>48</b> (2017)	<b>17</b> (2017)	<b>6.5</b> (2017)	<b>26</b> (2016)	<b>7.3</b> (2016)	<b>55</b> (2017)	<b>10.0</b> (2016)	<b>20</b> (2016)	<b>8.7</b> (2015)	<b>28</b> (2017)
Male (15-64) population (M)	<b>0.7</b> (2016)	<b>5.2</b> (2018)	<b>6.3</b> (2016)	<b>0.4</b> (2017)	<b>14</b> (2017)	<b>4.9</b> (2017)	<b>7.2</b> (2016)	<b>50</b> (2016)	<b>18</b> (2017)	<b>14</b> (2016)	<b>10</b> (2016)	<b>4.1</b> (2015)	<b>4.4</b> (2017)
Population (M)	<b>2.2</b> (2018)	<b>20</b> (2018)	<b>26</b> (2018)	<b>1.4</b> (2018)	<b>51</b> (2018)	<b>19</b> (2018)	<b>31</b> (2018)	<b>196</b> (2018)	<b>58</b> (2018)	<b>59</b> (2018)	<b>44</b> (2018)	<b>18</b> (2018)	<b>14</b> (2018)

SOURCE INFORMATION:

All condom use and knowledge data is from the DHS Stat Compiler with the following exceptions: Botswana (BAIS IV with combined data for men and women as presented in the summary report); South Africa (condom use indicator is % of men/women aged 15-49 who had more than one partner in the past 12 months reporting the use of a condom during their last sexual intercourse); MSM (UNAIDS 2017 Prevention Scorecard).

Condom distribution data: social marketing sector is from DKT's contraceptive statistics report; all other sectors from MGH reports from various unpublished sources.

Total population is drawn from World Bank data.

Adult male population is drawn from the reference tables of the UNAIDS Condom Needs Estimation Tool v 1.2.

HIV prevalence is drawn from Avert.org.

Empty cells indicate that data were not available, more than 10 years out of date, considered unreliable or otherwise not comparable to other countries...