South Africa		Based on most recent available data.	Versio	n 2.01.				
Output (coverage)		Outcome (service use/beha	aviour)		•	Impact		Summary
Condoms					New IIIV in	factions (acce 15)	in 1000a)	Scores
Number of condoms distributed and sold / year (in millions)	636	Condom use with non-regular partners (%)	Women 15-49	61	270	fections (ages 15+,	in 1000s)	AGYW&MPs
Number of condoms distributed and sold per man 15-64 /year	34	Condom use with non-regular partners (%)	Men 15-49	73	400 370			5
% of condom distribution need met for men 15-64	92				350			SW
Voluntary medical male circumcision (VMMC)					300			6
Number of VMMCs performed / year (in thousands)	452	% of 2020 VMMC target achieved	Men	50	250		190	MSM
% of annual VMMC target achieved	58	National male circumcision prevalence (%)	Men 15-24	70	200			7
		realistic file circums son prevalence (78)	Men 15-49	57	150		93	PWID
ARV-based prevention					100		•	2
Composite PrEP score (0-10)	5	% of national PrEP need met	All pop.	id	50			Condoms
Number of people who received PrEP at least once in the past 12 months	42255	% of PLHIV virally surpressed	Women 15+	58		2015	2020	8
% of PLHIV on ART	65	% of PLHIV virally surpressed	Men 15+	53	2010	2015	2020	VMMC
Key populations						HIV	prevalence	6
Sex workers (SW)								PrEP
Population size estimate for sex workers (in thousands)	125	Condom use / last paid sex (%)	Sex workers	86	Sex v	vorkers <25 years	id	5
% of SWs who received at least two HIV prevention interventions (past 3 mo)	45	Condom use / last paid sex (%)	Men 15-49	83	Se	x workers all ages	58	
Prevention strategy includes core elements of SW prevention package	All	% of PLHIV on ART	SWs LHIV	24				
Gay men and other men who have sex with men (MSM)								
Population size estimate for men who have sex with men (in thousands)	347	Condom use / last anal sex (%)	MSM	98		MSM <25 years	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	33	% of PLHIV on ART	MSM LHIV	28		MSM all ages	18	
Prevention strategy includes core elements of MSM prevention package	All							
People who inject drugs (PWID)								
Population size estimate for people who inject drugs (in thousands)	75	% with safe injecting practices	PWID	id		PWID <25 years	12	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	24	% on opioid substitution therapy	PWID	id		PWID all ages	22	
Prevention strategy includes core elements of PWID harm reduction package	All	% of PLHIV on ART	PWID LHIV	id				
Structural barriers and enablers								
Criminalization of selling sex	Yes	26.5	Sex workers	id				
Criminalization of same sex relations	No	% of people who avoided health care because of stigma and discrimination	MSM	id				
Criminalization of drug use/consumption or possession for personal use	Yes	because of stigma and discimination	PWID	id				
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								
% of priority locations/districts with dedicated programs for AGYW & partners	29	Condom use with non-regular partners (%)	YW 15-24	47	You	ing women 15-24	10.2	
% of girls who completed lower secondary education	91		YM 15-24	68		Young men 15-24	3.4	
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	18		Adults 15-49	19.0	
Laws requiring parental consent for adolescents to access SRH services	No	% of women experienced physical or sexual	YW 15-19	30				
Provider-initiated condom promotion integrated into SRH services	Yes	violence from husband/partner	Women 15-49	30				
HIV testing services integrated with SRH services	Yes							
Acronyms na not applicable;		Very good			Low			
PLHIVpeople living with HIV; LHIVliving with HIV;YWyoung women;YMyoung men;		Good			Very low			
OST opioid substitution therapy; SRH sexual and reproductive health.		Medium		id	Insufficient data			