

Select country from dropdown menu

Angola

AGO

Based on most recent available data.

Version 2.02.

2023

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact	
Condoms				New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)	
Number of condoms distributed and sold / year (in millions)	8	Condom use with non-regular partners (%)	Women 15-49	27	<p>2010 baseline 2020, 2022 2020 target 2025 target</p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>
Number of condoms distributed/sold per couple-year (age range 15-64)	1		Men 15-49	46	
Estimated condom distribution need met (%)	3				
Men and boys (including VMMC)					
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na	
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	96	
			Men 15-49	96	
ARV-based prevention					
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	id	
Number of people who received PrEP at least once in the past 12 months	id	% of PLHIV virally suppressed	Women 15+	id	
% of PLHIV on ART	46		Men 15+	id	
Key populations					
<i>Sex workers (SW)</i>					
Population size estimate for sex workers	id	Condom use / last paid sex (%)	Sex workers	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	51		Men 15-49	71	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	42	
<i>Gay men and other men who have sex with men (MSM)</i>					
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	3	% on ART	MSM LHIV	id	
Prevention strategy includes core elements of MSM prevention package	> Half				
<i>People who inject drugs (PWID)</i>					
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID LHIV	id	
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART			
<i>Structural barriers and enablers</i>					
Criminalization of sex work	No	% of people who avoided health care because of stigma and discrimination	Sex workers	id	
Criminalization of same-sex sexual acts	No		MSM	id	
Criminalization of drug use/consumption or possession for personal use	id		PWID	id	
Criminalization of transgender people	Yes		Transgender	id	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings					
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	31	
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	46	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	32	
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	id	
HIV testing services integrated with SRH services	Yes		Women 15-49	id	

Summary Scores

AGYW	3
Sex workers	6
MSM	id
PWID	id
Condoms	2
VMMC	na
ART	5
PrEP	id

Acronyms: na ... not applicable;
PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
SRH ... sexual and reproductive health.

Very good
Good
Medium

Low
Very low
Insufficient data