

Select country from dropdown menu

Angola

AGO

Based on most recent available data.

Version 2.02.

2024

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms		Condom use with non-regular partners (%)		New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)	
Number of condoms distributed and sold / year (in millions)	23	Women 15-49	27	2010 baseline	21 000
Number of condoms distributed/sold per couple-year (age range 15-64)	3	Men 15-49	46	2020, 2022	14 000
Estimated condom distribution need met (%)	12			2020 target	5 100
				2025 target	4 100
Men and boys (including VMMC)		% of 2020 VMMC target achieved		HIV prevalence	
Number of VMMCs performed / year (in thousands)	na	Men	na	2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.	
% of annual VMMC target achieved	na	Men 15-24	96		
		Men 15-49	96		
ARV-based prevention		% of national PrEP need met			
Composite PrEP score (0-10)	1	All pop.	0		
Number of people who received PrEP at least once in the past 12 months	0	Women 15+	id		
% of PLHIV on ART	50	Men 15+	id		
Key populations		% of PLHIV virally suppressed			
Sex workers (SW)					
Population size estimate for sex workers	id	Sex workers	id	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	51	Men 15-49	71	Sex workers all ages	id
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	id		
Gay men and other men who have sex with men (MSM)					
Population size estimate for men who have sex with men	id	MSM	id	MSM <25 years	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	3	MSM LHIV	id	MSM all ages	id
Prevention strategy includes core elements of MSM prevention package	> Half	% on ART	id		
People who inject drugs (PWID)					
Population size estimate for people who inject drugs	id	% with safe injecting practices	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	id		
Structural barriers and enablers					
Criminalization of sex work	No	% of people who avoided health care because of stigma and discrimination	id		
Criminalization of same-sex sexual acts	No		id		
Criminalization of drug use/consumption or possession for personal use	id		id		
Criminalization of transgender people	#N/A		id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings					
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	Women 15-24	0.7
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	Men 15-24	0.3
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16	% who completed lower secondary education	Girls	Adults, 15+	1.4
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19		
HIV testing services integrated with SRH services	Yes		Women 15-49		

Summary Scores	
AGYW	3
Sex workers	6
MSM	id
PWID	id
Condoms	2
VMMC	na
ART	5
PrEP	1

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data