

Select country from dropdown menu

Cameroon

CMR

Based on most recent available data.

Version 2.02.

2023

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact	
Condoms				New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)	
Number of condoms distributed and sold / year (in millions)	27	Condom use with non-regular partners (%)	Women 15-49	43	<p>2010 baseline</p> <p>2020, 2022</p> <p>2020 target</p> <p>2025 target</p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>
Number of condoms distributed/sold per couple-year (age range 15-64)	3		Men 15-49	63	
Estimated condom distribution need met (%)	16				
Men and boys (including VMMC)					
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na	
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	92	
			Men 15-49	94	
ARV-based prevention					
Composite PrEP score (0-10)	3	% of national PrEP need met	All pop.	40	
Number of people who received PrEP at least once in the past 12 months	4377	% of PLHIV virally suppressed	Women 15+	84	
% of PLHIV on ART	88		Men 15+	77	
Key populations					
Sex workers (SW)					
Population size estimate for sex workers	126400	Condom use / last paid sex (%)	Sex workers	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	8		Men 15-49	83	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	99	
Gay men and other men who have sex with men (MSM)					
Population size estimate for men who have sex with men	7000	Condom use / last anal sex (%)	MSM	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	10	% on ART	MSM LHIV	97	
Prevention strategy includes core elements of MSM prevention package	> Half				
People who inject drugs (PWID)					
Population size estimate for people who inject drugs	3500	% with safe injecting practices	PWID	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	26	% on opioid substitution therapy	PWID LHIV	id	
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	id	
Structural barriers and enablers					
Criminalization of sex work	Yes		Sex workers	5	
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of stigma and discrimination	MSM	id	
Criminalization of drug use/consumption or possession for personal use	No		PWID	id	
Criminalization of transgender people	No		Transgender	id	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings					
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	50	
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	65	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16	% who completed lower secondary education	Girls	43	
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	20	
HIV testing services integrated with SRH services	Yes		Women 15-49	22	

Summary Scores

AGYW	5
Sex workers	5
MSM	id
PWID	3
Condoms	3
VMMC	na
ART	9
PrEP	3

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data