Select country from dropdown menu	This scoreca	ard for Central African Republic was automatically created based	l on global databases an	id was not	quality asssured as part of the GPC's annual revie	w process.	
Central African Republic	CAF	Based on most recent available data.	Version	2.02.		2023	
out (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		
Number of condoms distributed and sold / year (in millions)	id	Condom use with non-regular partners (%)	Women 15-49	id	(ages 15+ trend vs. 2020 and 2025 targets)		2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	id		Men 15-49	id	9′000 _8 200	200	
Estimated condom distribution need met (%)	id				8'000 7 300	7 ′400	◆2020 target
Men and boys (including VMMC)					7'000	400	2025 target
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na	6'000		
% of annual VMMC target achieved na National male circum	National male sireumaisian provolence (0/)	Men 15-24	id	5'000		2020 and 20	
		National male circumcision prevalence (%)	Men 15-49	id	4′000		targets repr
ARV-based prevention					3′000 2′000 ♦	1 600	required contribution
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	id	1′000 2 100	•	global targe
Number of people who received PrEP at least once in the past 12 months	id	0/ of DILIN/virally average of	Women 15+	id			75% reducti 2020 and 82
% of PLHIV on ART	% of PLHIV on ART 49	Men 15+	id	2010 2015 2020	2025	reduction by	
Key populations					HIV	prevalence	2025 agains 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	3900	0 1 (1 1 1 10)	Sex workers	80	Sex workers <25 years	8.9	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	57	Condom use / last paid sex (%)	Men 15-49	id	Sex workers all ages	6.1	
Prevention strategy includes core elements of SW prevention package	< Half	% on ART	SWs LHIV	61			
Gay men and other men who have sex with men (MSM)							Summa
Population size estimate for men who have sex with men	3000	Condom use / last anal sex (%)	MSM	30	MSM <25 years	4.7	Score
% of MSM who received at least two HIV prevention interventions (past 3 mo)	42	% on ART	MSM LHIV	60	MSM all ages	3.4	AGYV
Prevention strategy includes core elements of MSM prevention package	> Half						1
People who inject drugs (PWID)							Sex worl
Population size estimate for people who inject drugs	id	% with safe injecting practices		id	PWID <25 years	id	6
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id			4
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	85			id
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	id			Condor
Criminalization of drug use/consumption or possession for personal use	Yes	 stigma and discrimination	PWID	id			id
Criminalization of transgender people	No		Transgender	id			VMM
dolescent girls, young women (AGYW) & partners in high-HIV incidence settings							na
% of priority locations/districts with dedicated programs for AGYW & partners	id		Women 15-24	id	Women 15-24	1.8	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	id	Men 15-24	0.9	5
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <18	% who completed lower secondary education	Girls	10	Adults, 15+	3.5	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id			id
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id			
ms: na not applicable;		Very good			Low		
people living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		