

Select country from dropdown menu

Central African Republic

This scorecard for Central African Republic was automatically created based on global databases and was not quality assured as part of the GPC's annual review process.

CAF

Based on most recent available data.

Version 2.02.

2024

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms			New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)			
Number of condoms distributed and sold / year (in millions)	id	Condom use with non-regular partners (%)	Women 15-49	id	<p>2010 baseline 2020, 2022 2020 target 2025 target</p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>	
Number of condoms distributed/sold per couple-year (age range 15-64)	id		Men 15-49	id		
Estimated condom distribution need met (%)	id					
Men and boys (including VMMC)						
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	id		
			Men 15-49	id		
ARV-based prevention						
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	id	% of PLHIV virally suppressed	Women 15+	id		
% of PLHIV on ART	id		Men 15+	id		
Key populations			HIV prevalence			
Sex workers (SW)						
Population size estimate for sex workers	id	Condom use / last paid sex (%)	Sex workers	80	Sex workers <25 years	8.9
% of SWs who received at least two HIV prevention interventions (past 3 mo)	57	% on ART	Men 15-49	id	Sex workers all ages	6.1
Prevention strategy includes core elements of SW prevention package	< Half		SWs LHIV	61		
Gay men and other men who have sex with men (MSM)						
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	30	MSM <25 years	4.7
% of MSM who received at least two HIV prevention interventions (past 3 mo)	42	% on ART	MSM LHIV	60	MSM all ages	3.4
Prevention strategy includes core elements of MSM prevention package	> Half					
People who inject drugs (PWID)						
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID LHIV	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART				
Structural barriers and enablers						
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	85		
Criminalization of same-sex sexual acts	No		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Criminalization of transgender people	No		Transgender	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	id	Women 15-24	id
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	id	Men 15-24	id
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	10	Adults, 15+	id
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	id		
HIV testing services integrated with SRH services	Yes		Women 15-49	id		

Summary Scores

AGYW	1
Sex workers	6
MSM	4
PWID	id
Condoms	id
VMMC	na
ART	id
PrEP	id

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data