Select country from dropdown menu	This scoreca	rd for Congo was automatically created based on global databa	ises and was not qualit	y asssured a	is part of the GPC's annual review process.		
Congo	COG	Based on most recent available data.	Versio	n 2.02 .		2023	
Dutput (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		2010 hasslins
Number of condoms distributed and sold / year (in millions)	id	Condem use with non-regular portners (0/)	Women 15-49 40 Men 15-49 64		(ages 15+ trend vs. 2020 and 2025 targets)		2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	id	Condom use with non-regular partners (%)			14'000	13'000	• 2020, 2022
Estimated condom distribution need met (%)	id				12'000 11 00		2020 target
Men and boys (including VMMC)					10'000		2025 target
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na			
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	98	8'000 6 100		2020 and 2025 targets represe
			Men 15-49	99	6'000		the country's
ARV-based prevention					4'000	4 200	required contribution to
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	0	2'000	1 200	global targets,
Number of people who received PrEP at least once in the past 12 months	id	% of PLHIV virally surpressed	Women 15+	id	1 500	•	75% reduction 2020 and 82.5
% of PLHIV on ART	24		Men 15+	id	2010 2015 2020	2025	reduction by
Key populations					н	V prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	id		Sex workers	86	Sex workers <25 years	3.5	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	28	Condom use / last paid sex (%)	Men 15-49	75	Sex workers all ages	8.1	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id			
Gay men and other men who have sex with men (MSM)							Summary
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	64	MSM <25 years	32.2	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	67	% on ART	MSM LHIV	id	MSM all ages	41.2	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						4
People who inject drugs (PWID)							Sex worke
Population size estimate for people who inject drugs	id	% with safe injecting practices		id	PWID <25 years	id	5
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id			7
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	17			id
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	10			Condom
Criminalization of drug use/consumption or possession for personal use	id	stigma and discrimination	PWID	id			id
Criminalization of transgender people	No		Transgender	id			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings			-				na
% of priority locations/districts with dedicated programs for AGYW & partners	id		Women 15-24	45	Women 15-24	3.6	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	66	Men 15-24	0.7	2
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	45	Adults, 15+	4.0	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id			id
HIV testing services integrated with SRH services	Partial	violence from husband/partner	Women 15-49	id			
cronyms: na not applicable;		Very good			Low		
HIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		
RH sexual and reproductive health.		Medium		id	Insufficient data		