Select country from dropdown menu							
Cote d'Ivoire	CIV	Based on most recent available data.	Version	2.02.		2023	
Output (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		2010 haveling
Number of condoms distributed and sold / year (in millions)	id	Condom use with non-regular partners (%)	Women 15-49 38		(ages 15+ trend vs. 2020 and 2025 targets)		2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	id	Condom use with non-regular partners (%)	Men 15-49	63	25'000		• 2020, 2022
Estimated condom distribution need met (%)	id				_20 000		2020 target
Men and boys (including VMMC)					20'000		2025 target
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na	15'000		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	96			2020 and 2025 targets represent
			Men 15-49	97	10'000 8 300)	the country's
ARV-based prevention					5/000	7'300 4 000	required contribution to
Composite PrEP score (0-10)	5	% of national PrEP need met	All pop.	40	5'000 5 000	•	global targets, a
Number of people who received PrEP at least once in the past 12 months	676	0/ of DUUN/virally ourproceed	Women 15+	69			75% reduction by 2020 and 82.5%
% of PLHIV on ART	72	% of PLHIV virally surpressed	Men 15+	54	2010 2015 2020	2025	reduction by
Key populations					н	V prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	66600	Condom use / last paid sex (%)	Sex workers	92	Sex workers <25 years	1.7	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	23		Men 15-49	63	Sex workers all ages	4.8	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id			
Gay men and other men who have sex with men (MSM)							Summary
Population size estimate for men who have sex with men	56000	Condom use / last anal sex (%)	MSM	75	MSM <25 years	5.5	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	39	% on ART	MSM LHIV	id	MSM all ages	7.7	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						na
People who inject drugs (PWID)							Sex workers
Population size estimate for people who inject drugs	2600	% with safe injecting practices	PWID	id	PWID <25 years	id	5
% of PWID who received at least two HIV prevention interventions (past 3 mo)	85	% on opioid substitution therapy	PWID	id	PWID all ages	3.4	MSM
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	id			6
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	14			9
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	6			Condoms
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	9			id
Criminalization of transgender people	No		Transgender	23			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							na
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom uses with non-requirementary (a)	Women 15-24	42	Women 15-24	0.8	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	67	Men 15-24	0.5	7
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16	% who completed lower secondary education	Girls	22	Adults, 15+	2.2	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id			5
HIV testing services integrated with SRH services	id	violence from husband/partner	Women 15-49	id			
Acronyms: na not applicable;		Very good			Low		
PLHIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good		: 4	Very low		
SRH sexual and reproductive health.		Medium		id	Insufficient data		