Select country from dropdown menu							
Democratic Republic of the Congo		Based on most recent available data.	Version 2.02.		2023		
utput (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		
Number of condoms distributed and sold / year (in millions)	3	Condom use with non-regular partners (%)	Women 15-49	24	(ages 15+ trend vs. 2020 and 2025	targets)	2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	0	Condom use with non-regular partners (%)	Men 15-49	33	30'000		• 2020, 2022
Estimated condom distribution need met (%)	1				25 000		2020 target
Men and boys (including VMMC)							2025 target
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na	20'000		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	id	15'000	000	2020 and 202 targets repre
			Men 15-49	97	10'000	0/000	the country's
ARV-based prevention					10 000	8'60g 100	required contribution
Composite PrEP score (0-10)	3	% of national PrEP need met	All pop.	12	5'000 6 40	00	global target
Number of people who received PrEP at least once in the past 12 months	8650	% of PLHIV virally surpressed	Women 15+	id	·		75% reductio 2020 and 82
% of PLHIV on ART	82	% OF PLHTV VIrally surpressed	Men 15+	78	2010 2015 20	20 2025	reduction by
Key populations						HIV prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	526000	Condom use / last paid sex (%)	Sex workers	48	Sex workers <25 yea	ars 4.8	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	38		Men 15-49	34	Sex workers all ag	es 7.5	
Prevention strategy includes core elements of SW prevention package	All	% on ART	SWs LHIV	id			
Gay men and other men who have sex with men (MSM)							Summa
Population size estimate for men who have sex with men	194900	Condom use / last anal sex (%)	MSM	57	MSM <25 yea	ars id	Score
% of MSM who received at least two HIV prevention interventions (past 3 mo)	39	% on ART	MSM LHIV	id	MSM all ag	es 7.1	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						4
People who inject drugs (PWID)							Sex work
Population size estimate for people who inject drugs	168000	% with safe injecting practices		23	PWID <25 yea	ars 0.6	3
% of PWID who received at least two HIV prevention interventions (past 3 mo)	23	% on opioid substitution therapy	PWID	id	PWID all ag	es 3.9	MSM
Prevention strategy includes core elements of PWID harm reduction package	#N/A	% on ART	PWID LHIV	id			5
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	id			1
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	18			Condor
Criminalization of drug use/consumption or possession for personal use	No	stigma and discrimination	PWID	14			1
Criminalization of transgender people	No		Transgender	18			VMM
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							na
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-results sectors (0)	Women 15-24	25	Women 15-	24 0.2	ART
Educational policies on HIV & sexuality education (secondary school)	id	Condom use with non-regular partners (%)	Men 15-24	33	Men 15-	24 0.2	8
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <18	% who completed lower secondary education	Girls	52	Adults, 1	5+ 0.6	PrEP
Provider-initiated condom promotion integrated into SRH services	No/id	% of women experienced physical or sexual	Women 15-19	id			3
HIV testing services integrated with SRH services	#N/A	violence from husband/partner	Women 15-49	id			
nyms: na not applicable;		Very good			Low		
people living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		
sexual and reproductive health.		Medium		id	Insufficient data		