

Select country from dropdown menu

Ethiopia

ETH

Based on most recent available data.

Version 2.02.

2024

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms		Condom use with non-regular partners (%)		New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)	
Number of condoms distributed and sold / year (in millions)	71	Women 15-49	20		
Number of condoms distributed/sold per couple-year (age range 15-64)	2	Men 15-49	51		
Estimated condom distribution need met (%)	27				
Men and boys (including VMMC)		% of 2020 VMMC target achieved			
Number of VMMCs performed / year (in thousands)	27	Men	na		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)			
		Men 15-24	88		
		Men 15-49	91		
ARV-based prevention		% of national PrEP need met			
Composite PrEP score (0-10)	5	All pop.	28		
Number of people who received PrEP at least once in the past 12 months	16055	% of PLHIV virally suppressed			
% of PLHIV on ART	85	Women 15+	77		
		Men 15+	77		
Key populations				HIV prevalence	
Sex workers (SW)				Sex workers <25 years	id
Population size estimate for sex workers	id	Condom use / last paid sex (%)	Sex workers	Sex workers all ages	18.3
% of SWs who received at least two HIV prevention interventions (past 3 mo)	79		Men 15-49		
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV		
			id		
Gay men and other men who have sex with men (MSM)				MSM <25 years	id
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	MSM all ages	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id	% on ART	MSM LHIV		
Prevention strategy includes core elements of MSM prevention package	id		id		
			id		
People who inject drugs (PWID)				PWID <25 years	id
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	PWID all ages	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	id		
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV		
			id		
Structural barriers and enablers					
Criminalization of sex work	Yes		Sex workers		
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of stigma and discrimination	MSM		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID		
Criminalization of transgender people	No		Transgender		
			id		
			id		
			id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings					
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	Women 15-24	0.3
Educational policies on HIV & sexuality education (secondary school)	No		Men 15-24	Men 15-24	0.2
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16	% who completed lower secondary education	Girls	Adults, 15+	0.9
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19		
HIV testing services integrated with SRH services	Partial		Women 15-49		
			id		
			id		

2010 baseline  
2020, 2022  
2020 target  
2025 target

2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

Summary Scores	
AGYW	2
Sex workers	8
MSM	id
PWID	id
Condoms	3
VMMC	na
ART	8
PrEP	5

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good  
Good  
Medium

Low  
Very low  
Insufficient data