Select country from dropdown menu							
Malawi	MWI	Based on most recent available data.	Version	2.02.		2023	
Output (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infection	ons	
Number of condoms distributed and sold / year (in millions)	120	Condom use with non-regular partners (%)	Women 15-49	49	(ages 15+ trend vs. 2020 and 2	025 targets)	2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	22		Men 15-49 73		45'000		2020, 2022
Estimated condom distribution need met (%)	100				40 000		◆2020 target
Men and boys (including VMMC)					35'000		2025 target
Number of VMMCs performed / year (in thousands)	259	% of 2020 VMMC target achieved	Men	11	30'000		
% of annual VMMC target achieved	30	National male circumcision prevalence (%)	Men 15-24 Men 15-49	29 28	20 000	18 000	2020 and 2025 targets represer the country's
ARV-based prevention					15'000	13′089	00 required
Composite PrEP score (0-10)	5	% of national PrEP need met	All pop.	53	10'000 5'000 1	10 000	contribution to global targets, a
Number of people who received PrEP at least once in the past 12 months	23104		Women 15+	93			75% reduction be 2020 and 82.5%
% of PLHIV on ART	93	% of PLHIV virally surpressed	Men 15+	81	2010 2015	2020 202	
Key populations						HIV prevalen	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	39000	0 1 1 1 10	Sex workers	65	Sex workers <25	years id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	68	Condom use / last paid sex (%)	Men 15-49	75	Sex workers all	ages 49.9	
Prevention strategy includes core elements of SW prevention package	< Half	% on ART	SWs LHIV	90			
Gay men and other men who have sex with men (MSM)							Summary
Population size estimate for men who have sex with men	50000	Condom use / last anal sex (%)	MSM	79	MSM <25 y	years 4.6	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	65	% on ART	MSM LHIV	93	MSM all	ages 12.9	AGYW
Prevention strategy includes core elements of MSM prevention package	< Half						4
People who inject drugs (PWID)							Sex worke
Population size estimate for people who inject drugs	8400	% with safe injecting practices	DIAME	id	PWID <25	years id	6
% of PWID who received at least two HIV prevention interventions (past 3 mo)	0	% on opioid substitution therapy	PWID	id	PWID all	ages id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id			7
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	49			0
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of	MSM	13			Condoms
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id			8
Criminalization of transgender people	Yes		Transgender	id			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							2
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (n/)	Women 15-24	53	Women 1	15-24 2.2	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	73	Men 1	1.5	9
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	21	Adults	, 15+ 8.0	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id			5
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id			
Acronyms: na not applicable;		Very good			Low		
PLHIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good		: 4	Very low		
RH sexual and reproductive health.		Medium		Ia	Insufficient data		