

Select country from dropdown menu

Mexico

MEX

Based on most recent available data.

Version 2.02.

2023


Output (coverage)

Outcome (service use/behaviour)

Impact

New HIV infections

(ages 15+ trend vs. 2020 and 2025 targets)



2010 baseline
2020, 2022
2020 target
2025 target

2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

ARV-based prevention

| | |
|--|------|
| Composite PrEP score (0-10) | 4 |
| Number of people who received PrEP at least once in the past 12 months | 7594 |
| % of PLHIV on ART | 62 |

| | | |
|-------------------------------|-----------|----|
| % of national PrEP need met | All pop. | 3 |
| % of PLHIV virally suppressed | Women 15+ | 51 |
| | Men 15+ | 59 |

Key populations

Sex workers (SW)

| | |
|---|--------|
| Population size estimate for sex workers | 240000 |
| % of SWs who received at least two HIV prevention interventions (past 3 mo) | 0 |
| Prevention strategy includes core elements of SW prevention package | > Half |

| | | |
|--------------------------------|-------------|----|
| Condom use / last paid sex (%) | Sex workers | id |
| % on ART | Men 15-49 | id |
| | SWs LHIV | id |

| | |
|-----------------------|-----|
| Sex workers <25 years | id |
| Sex workers all ages | 0.8 |

Gay men and other men who have sex with men (MSM)

| | |
|---|---------|
| Population size estimate for men who have sex with men | 1200000 |
| % of MSM who received at least two HIV prevention interventions (past 3 mo) | 0 |
| Prevention strategy includes core elements of MSM prevention package | > Half |

| | | |
|--------------------------------|----------|----|
| Condom use / last anal sex (%) | MSM | id |
| % on ART | MSM LHIV | id |

| | |
|---------------|------|
| MSM <25 years | id |
| MSM all ages | 11.9 |

People who inject drugs (PWID)

| | |
|--|-----|
| Population size estimate for people who inject drugs | id |
| % of PWID who received at least two HIV prevention interventions (past 3 mo) | id |
| Prevention strategy includes core elements of PWID harm reduction package | All |

| | | |
|----------------------------------|-----------|----|
| % with safe injecting practices | PWID | id |
| % on opioid substitution therapy | PWID | 9 |
| % on ART | PWID LHIV | id |

| | |
|----------------|-----|
| PWID <25 years | id |
| PWID all ages | 2.9 |

Structural barriers and enablers

| | |
|--|----|
| Criminalization of sex work | No |
| Criminalization of same-sex sexual acts | No |
| Criminalization of drug use/consumption or possession for personal use | No |
| Criminalization of transgender people | No |

| | | |
|--|-------------|----|
| % of people who avoided health care because of stigma and discrimination | Sex workers | id |
| | MSM | id |
| | PWID | id |
| | Transgender | id |

| | |
|-------------|-----|
| Women 15-24 | 0.1 |
| Men 15-24 | 0.3 |
| Adults, 15+ | 0.4 |

Summary Scores

| | |
|-------------|----|
| Sex workers | id |
| MSM | id |
| PWID | 0 |
| ART | 6 |
| PrEP | 4 |

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

| |
|-----------|
| Very good |
| Good |
| Medium |

| |
|-------------------|
| Low |
| Very low |
| Insufficient data |