Select country from dropdown menu								
Mozambique	MOZ	Based on most recent available data.	Version	n 2.02.			202	3
Output (coverage)		Outcome (service use/behaviour)				Impact		
Condoms					1			
Number of condoms distributed and sold / year (in millions)	id	6 1 11 10 10 100	Women 15-49	42	(ages 15+ trend vs. 2020 and 2025 to		025 targets)	2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	id	Condom use with non-regular partners (%)	Men 15-49	47	160'000			2020, 2022
Estimated condom distribution need met (%)	id				140'000 130	000		◆2020 target
Men and boys (including VMMC)					120'000		110 000	2025 target
Number of VMMCs performed / year (in thousands)	172	% of 2020 VMMC target achieved	Men	27	100'000			
% of annual VMMC target achieved	77	National male size maising providence (0/)	Men 15-24	66	80'000		84'00	
	National male circumcision prevalence (%)	Men 15-49	47	60'000			targets represe the country's	
ARV-based prevention					40'000		26	000 required contribution to
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	id	20'000	3	32 000	global targets,
Number of people who received PrEP at least once in the past 12 months	id	or 500000 to 11	Women 15+	74				75% reduction 2020 and 82.59
% of PLHIV on ART	81	% of PLHIV virally surpressed	Men 15+	70	2010	2015	2020 2	025 reduction by
Key populations							HIV prevale	2025 against 2010 as a
Sex workers (SW)								baseline.
Population size estimate for sex workers	224000		Sex workers	id		Sex workers <25 y	years id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	57	Condom use / last paid sex (%)	Men 15-49	31		Sex workers all	ages id	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id				
Gay men and other men who have sex with men (MSM)								Summary
Population size estimate for men who have sex with men	64000	Condom use / last anal sex (%)	MSM	id		MSM <25 y	years id	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	31	% on ART	MSM LHIV	id		MSM all	ages id	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half							3
People who inject drugs (PWID)								Sex worke
Population size estimate for people who inject drugs	14000	% with safe injecting practices		id		PWID <25 y	years id	4
% of PWID who received at least two HIV prevention interventions (past 3 mo)	40	% on opioid substitution therapy	PWID	id		PWID all	ages id	MSM
Prevention strategy includes core elements of PWID harm reduction package	< Half	% on ART	PWID LHIV	id				id
Structural barriers and enablers								PWID
Criminalization of sex work	No		Sex workers	id				4
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	id				Condom
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id				id
Criminalization of transgender people	No		Transgender	id				VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								5
% of priority locations/districts with dedicated programs for AGYW & partners	33		Women 15-24	51		Women 1	5-24 6.5	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	48		Men 1		8
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	11		Adults		PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id				id
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id				
cronyms: na not applicable;		Very good			Low			
LHIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low			
RH sexual and reproductive health.		Medium		id	Insufficient data			