2.02.	2.02.	Version	Based on most recent available data.	MOZ	Mozambique
	$\rightarrow$		Outcome (service use/behaviour)		Dutput (coverage)
New HIV					Condoms
37 (ages 15+ trend vs. 2	37		Condom use with non-regular partners (%)	109	Number of condoms distributed and sold / year (in millions)
<b>47</b> 140,000 <u>130 000</u>	47			12	Number of condoms distributed/sold per couple-year (age range 15-64)
120,000				48	Estimated condom distribution need met (%)
					Men and boys (including VMMC)
<b>27</b> 100,000	27	Men	% of 2020 VMMC target achieved	163	Number of VMMCs performed / year (in thousands)
id 80,000	id	Men 15-24	National male circumcision prevalence (%)	77	% of annual VMMC target achieved
75 60,000	75	Men 15-49			
40,000					ARV-based prevention
0 20,000	0	All pop.	% of national PrEP need met	3	Composite PrEP score (0-10)
82	82	Women 15+		0	Number of people who received PrEP at least once in the past 12 months
<b>72</b> 2010 2015	72	Men 15+	% of PLHIV virally surpressed	86	% of PLHIV on ART
					Key populations
					Sex workers (SW)
id Sex worke	id	Sex workers	Condom use / last paid sex (%)	id	Population size estimate for sex workers
id Sex wo	id	Men 15-49		57	% of SWs who received at least two HIV prevention interventions (past 3 mo)
id	id	SWs LHIV	% on ART	> Half	Prevention strategy includes core elements of SW prevention package
			·		Gay men and other men who have sex with men (MSM)
id MS	id	MSM	Condom use / last anal sex (%)	id	Population size estimate for men who have sex with men
id	id	MSM LHIV	% on ART	31	% of MSM who received at least two HIV prevention interventions (past 3 mo)
				> Half	Prevention strategy includes core elements of MSM prevention package
			•		People who inject drugs (PWID)
id PW	id		% with safe injecting practices	id	Population size estimate for people who inject drugs
id F		PWID	% on opioid substitution therapy	40	% of PWID who received at least two HIV prevention interventions (past 3 mo)
id	id	PWID LHIV	% on ART	< Half	Prevention strategy includes core elements of PWID harm reduction package
					Structural barriers and enablers
id	id	Sex workers		No	Criminalization of sex work
id	id	MSM	% of people who avoided health care because of	No	Criminalization of same-sex sexual acts
id	id	PWID	stigma and discrimination	Yes	Criminalization of drug use/consumption or possession for personal use
id	id	Transgender		No	Criminalization of transgender people
					Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings
41 W		Women 15-24	Condom use with non-regular partners (%)	35	% of priority locations/districts with dedicated programs for AGYW & partners
51		Men 15-24		Yes	Educational policies on HIV & sexuality education (secondary school)
11	11	Girls	% who completed lower secondary education	Yes, <12	Laws requiring parental consent for adolescents to access HIV testing services
id	id	Women 15-19	% of women experienced physical or sexual	Yes	Provider-initiated condom promotion integrated into SRH services
id	id	Women 15-49	violence from husband/partner	Yes	HIV testing services integrated with SRH services
Low			Very good		cronyms: na not applicable;
Very low			Good		HIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;
id Insufficient data	id		Medium		RH sexual and reproductive health.

