Select country from dropdown menu							
Namibia	NAM	Based on most recent available data.	Version	n 2.02.		2023	
Output (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		
Number of condoms distributed and sold / year (in millions)	11		Women 15-49 66		(ages 15+ trend vs. 2020 and 2025 targets)		2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	15	Condom use with non-regular partners (%)	Men 15-49	82	10'000 _9 300		• 2020, 2022
Estimated condom distribution need met (%)	46				9'000		2020 target
Men and boys (including VMMC)					8'000		2025 target
Number of VMMCs performed / year (in thousands)	21	% of 2020 VMMC target achieved	Men	20	7'000 6'000	5 800	
% of annual VMMC target achieved	59	National male sincumsision provalence (%)	Men 15-24	22	5'200		2020 and 2025
	National male circumcision prevalence (9	Men 15-49	26	4'000		targets represent the country's	
ARV-based prevention					3'000	1 900	required contribution to
Composite PrEP score (0-10)	10	% of national PrEP need met	All pop.	id	2'000 2 1'000 2	300	global targets, a
Number of people who received PrEP at least once in the past 12 months	29826		Women 15+	90			75% reduction k 2020 and 82.5%
% of PLHIV on ART	91	% of PLHIV virally surpressed	Men 15+	81	2010 2015	2020 2025	reduction by
Key populations						HIV prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	8500	Condom use / last paid sex (%)	Sex workers	42	Sex workers <25	ears 14.0	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	id		Men 15-49	67	Sex workers all	ages 29.9	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id			
Gay men and other men who have sex with men (MSM)							Summary
Population size estimate for men who have sex with men	2200	Condom use / last anal sex (%)	MSM	55	MSM <25	ears 3.6	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	33	% on ART	MSM LHIV	id	MSM all	ages 7.8	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						5
People who inject drugs (PWID)							Sex worker
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	id	PWID <25	ears id	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all	ages id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id			4
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	id			id
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of	MSM	id			Condoms
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id			6
Criminalization of transgender people	No		Transgender	id			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							4
% of priority locations/districts with dedicated programs for AGYW & partners	42	Condom use with non-regular partners (%)	Women 15-24	68	Women 1	5-24 5.7	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	condon use with non regular partners (70)	Men 15-24	84	Men 1	5-24 2.5	9
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	62	Adults	15+ 12.6	PrEP
Provider-initiated condom promotion integrated into SRH services	No/id	% of women experienced physical or sexual	Women 15-19	id			10
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id			
cronyms: na not applicable;		Very good			Low		
HIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		
RH sexual and reproductive health.		Medium		id	Insufficient data		