


The State of HIV Prevention in Nigeria


2023

Number of new HIV infections (all ages)




2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

Change in new HIV infections



HIV prevalence



HIV programme coverage and outcomes

ADOLESCENT GIRLS, YOUNG WOMEN & MALE PARTNERS

Condom use with a non-regular partner among young people 15-24 years old (%)

Young women: 38, Young men: 62

Target: 90%

Score: 5

KEY POPULATIONS

Sex workers

Condom use at last paid sex is (%)

Sex workers: 86, Clients: 74

Target: 95%

Score: 8

Gay men and other men who have sex with men

Condom use at last anal sex (%)

Gay men and other men who have sex with men: 70

Target: 90%

Score: 8

People who inject drugs

Use of harm reduction services (%)

With safe injections: 36, On opioid substitution therapy: 65

Target: 50%

Score: 2

CONDOM PROGRAMMING

Condom use with a non-regular partner, 15-49 years (%)

women: 36, men: 65

Target: 90%

Score: 6

MEN AND BOYS (INCLUDING VMMC)

Uptake of voluntary medical male circumcision

96%

Target: 90%

Score: na

ANTIRETROVIRAL DRUG-BASED PREVENTION

Pre-exposure prophylaxis

Number of people actively taking Pre-Exposure Prophylaxis (PrEP)

2017: 360, 2018: 118, 2019: 11413, 2020: 37736, 2021: 148952, 2022: 384429

Change in use of PrEP (2021-2022): +295%

Score: id

Antiretroviral treatment

People living with HIV virally suppressed (%)

Target: 86%

Score: id

PREVENTION OF VERTICAL TRANSMISSION OF HIV

MTCT rate

Score: id

% of HIV-positive pregnant women receiving ART

Score: id

Number of Estimated births to women living with HIV

Score: id

Number of new child infections due to vertical transmission

Score: id

Distribution of new child infections

- % of child infections because mother acquired HIV during pregnancy or breastfeeding
- % of child infections because mother did not receive antiretroviral therapy during pregnancy or breastfeeding
- % of child infections because mother did not continue antiretroviral treatment during pregnancy or breastfeeding
- % of child infections because mother was on antiretroviral treatment during pregnancy or breastfeeding, but was not virologically suppressed

Scores (1-10) Very good Good Medium Low Very low id ... insufficient data na ... not applicable

Policy and structural barriers

Key populations

	Sex workers	Gay men and other men who have sex with men	People who inject drugs	Transgender people	Prisoners
Criminalization of the behaviour of key populations	Yes	Yes	Yes	No	na
The national strategy includes critical elements of the programme package for key populations	> Half	> Half	> Half	< Half	> Half
Avoided health care because of stigma and discrimination	id	id	id	id	na
Population size	906 900	240 000	446 000	69 000	78 000

Adolescent girls and young women

	15-19 years	15-49 years
Proportion of women who experienced intimate partner violence	14%	13%
Girls who completed lower secondary education	59%	
Policies on life skills-based HIV and sexuality education (secondary schools)	Yes	
Laws requiring parental consent for adolescents to access HIV testing services, age of consent	Yes, <18	

Baseline status of 10 HIV Prevention 2025 Road Map Actions

Road Map Action	2023
1. Data-driven needs assessment	id
2. Precision prevention approach	id
3. Define investment needs	id
4. HIV prevention leadership agencies	id
5. Expand community-led services	id
6. Remove social and legal barriers	id
7. Integration with related services	id
8. Introduction of new technologies	id
9. Real-time programme monitoring	id
10. Accountability for HIV progress	id

Note: 'Yes' refers to the adaptation having been introduced (not necessarily it being universally available).

Linkages between HIV and sexual and reproductive health services

HIV testing services integrated within sexual and reproductive health	Yes
Provider-initiated condom promotion integrated into sexual and reproductive health services	Yes

Data sources: UNAIDS 2021 epidemiological estimates; Global AIDS Monitoring 2021; and ICF - the DHS Program STATcompiler. Data sources for key population program coverage: Global AIDS Monitoring 2022, Global Fund and PEPFAR reports obtained in 2022. Note: The 2023 UNAIDS epidemiological estimates represent the year 2022. Other data points may refer to various years when the surveys were conducted. Note: 2020 and 2025 targets for reducing new HIV infections represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline. These reductions are required to achieve a 90% reduction by 2030.