| Select country from dropdown menu | | | | | | | |
|--|-----------|--|-----------------------|----------|---|------------|---------------------------------------|
| Pakistan | PAK | Based on most recent available data. | Version | 2.02. | | 2023 | |
| Output (coverage) | | Outcome (service use/behaviour) | | | Impact | | |
| | | | | | New HIV infections (ages 15+ trend vs. 2020 and 2025 target | ets) | 2010 baseline |
| | | | | | 1 | | 2 020, 2022 |
| | | | | | 1 | | ◆2020 target |
| | | | | | 1 | | 2025 target |
| | | | | | 1 | | 2020 and 2025 |
| | | | | | 1 | | targets represent |
| ADV based provention | | | | | 0 | | the country's required |
| ARV-based prevention | : d | 0/ of national DrED nood mat | Allman | ; al | 0 | | contribution to |
| Composite PrEP score (0-10) Number of people who received PrEP at least once in the past 12 months | id id | % of national PrEP need met | All pop. Women 15+ | id id | 0 _0 - | 0 | global targets, a 75% reduction by |
| % of PLHIV on ART | 13 | % of PLHIV virally surpressed | Men 15+ | id | 2010 2015 02020 | 2025 | 2020 and 82.5% reduction by |
| Key populations | 13 | | Wich 15 | Id | HIV | prevalence | 2025 against 2010 |
| Sex workers (SW) | | | | | | • | as a baseline. |
| Population size estimate for sex workers | id | | Sex workers | id | Sex workers <25 years | id | |
| % of SWs who received at least two HIV prevention interventions (past 3 mo) | 6 | Condom use / last paid sex (%) | Men 15-49 | id | Sex workers all ages | id | |
| Prevention strategy includes core elements of SW prevention package | < Half | % on ART | SWs LHIV | 11 | | | |
| Gay men and other men who have sex with men (MSM) | | | | | | | |
| Population size estimate for men who have sex with men | id | Condom use / last anal sex (%) | MSM | id | MSM <25 years | id | |
| % of MSM who received at least two HIV prevention interventions (past 3 mo) | 6 | % on ART | MSM LHIV | 3 | MSM all ages | id | |
| Prevention strategy includes core elements of MSM prevention package | > Half | | | | | | |
| People who inject drugs (PWID) | | | | | | | Summary |
| Population size estimate for people who inject drugs | id | % with safe injecting practices | PWID | id | PWID <25 years | id | Scores |
| % of PWID who received at least two HIV prevention interventions (past 3 mo) | 37 | % on opioid substitution therapy | | id | PWID all ages | id | Sex workers |
| Prevention strategy includes core elements of PWID harm reduction package | < Half | % on ART | PWID LHIV | 29 | | | id |
| Structural barriers and enablers | V | | Carrantantan | : -1 | | | MSM |
| Criminalization of sex work | Yes | ov 6 | Sex workers | id | | | id |
| Criminalization of same-sex sexual acts | Yes | % of people who avoided health care because of stigma and discrimination | MSM PWID | id :d | | | PWID |
| Criminalization of drug use/consumption or possession for personal use Criminalization of transgender people | Yes No | 0.016.00 | Transgender | id id | | | ART |
| Chillinalization of transgender people | NO | | Hansgender | Iu | | | 1 |
| | | | | | Women 15-24 | 0.0 | PrEP |
| | | | | | Men 15-24 | 0.1 | id |
| | | | | | Adults, 15+ | 0.2 | |
| | | | | | · | | |
| | | | | | | | |
| Acronyms: na not applicable; | | Very good | | | Low | | |
| PLHIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men; | | Good | | | Very low | | |
| SRH sexual and reproductive health. | | Medium | | id | Insufficient data | | |