| Select country from dropdown menu | | | | | 2. 2024 | | | | |
|--|--|--|-------------|-------|---|-------------|--------------|---------------------------------|---------------------------------|
| | | | | | | | | | |
| Output (coverage) | | Outcome (service use/behaviour) | | | | Impa | | | |
| Condoms | | | | | New HIV infections | | | | 2010 baseline |
| Number of condoms distributed and sold / year (in millions) | 22 | Condom use with non-regular partners (%) | Women 15-49 | 46 | (ages 15+ trend vs. 2020 and 2025 targets | | | ets) | |
| Number of condoms distributed/sold per couple-year (age range 15-64) | 6 | 3 1 | Men 15-49 | 75 | 10,000 | 8 900 | | | 2020, 2022 |
| Estimated condom distribution need met (%) | 45 | | | | 9,000 8,000 | | | | 2020 target |
| Men and boys (including VMMC) | | | | | 7,000 | | | | 2025 target |
| Number of VMMCs performed / year (in thousands) | 240 | % of 2020 VMMC target achieved | Men | 94 | 6,000 | | | | 2020 1202 |
| % of annual VMMC target achieved | National male circumcision prevalence (9 | Men 15-24 | 73 | 5,000 | | | | 2020 and 2025 targets repres | |
| | | | Men 15-49 | 56 | 4,000 | | 2 800 | 2,700 | the country's required |
| ARV-based prevention | | | | | 3,000 2,000 | | | 1800 | contribution t |
| Composite PrEP score (0-10) | 10 | % of national PrEP need met | All pop. | id | 1,000 | | 2 200 | • | global targets 75% reduction |
| Number of people who received PrEP at least once in the past 12 months | 18582 | % of PLHIV virally surpressed | Women 15+ | 94 | | | | | 2020 and 82.5 |
| % of PLHIV on ART | % of PLHIV on ART 96 | | Men 15+ | 96 | 2 | 010 2015 | 2020 | 2025 | reduction by 2 against 2010 a |
| Key populations | | | | | | | HIV | prevalence | |
| Sex workers (SW) | | | | | | | | | |
| Population size estimate for sex workers | 98587 | Condom use / last paid sex (%) | Sex workers | 81 | | Sex workers | s <25 years | 4.7 | |
| % of SWs who received at least two HIV prevention interventions (past 3 mo) | id | Condom use / last paid sex (70) | Men 15-49 | 75 | | Sex work | ers all ages | 35.2 | |
| Prevention strategy includes core elements of SW prevention package | > Half | % on ART | SWs LHIV | 68 | | | | | |
| Gay men and other men who have sex with men (MSM) | | | | | | | | | Summa |
| Population size estimate for men who have sex with men | 28958 | Condom use / last anal sex (%) | MSM | 34 | | MSN | 1 <25 years | 2.7 | Scores |
| % of MSM who received at least two HIV prevention interventions (past 3 mo) | id | % on ART | MSM LHIV | 66 | | M | SM all ages | 5.8 | AGYW |
| Prevention strategy includes core elements of MSM prevention package | > Half | | | | | | | | 4 |
| People who inject drugs (PWID) | | | | | | | | | Sex work |
| Population size estimate for people who inject drugs | id | % with safe injecting practices | PWID | id | | PWIE | 0 <25 years | id | id |
| % of PWID who received at least two HIV prevention interventions (past 3 mo) | id | % on opioid substitution therapy | PVVID | id | | PW | /ID all ages | id | MSM |
| Prevention strategy includes core elements of PWID harm reduction package | None | % on ART | PWID LHIV | id | | | | | id |
| Structural barriers and enablers | | | | | | | | | PWID |
| Criminalization of sex work | Yes | | Sex workers | 11 | | | | | id |
| Criminalization of same-sex sexual acts | No | % of people who avoided health care because of | MSM | 4 | | | | | Condon |
| Criminalization of drug use/consumption or possession for personal use | id | stigma and discrimination | PWID | id | | | | | 5 |
| Criminalization of transgender people | No | | Transgender | id | | | | | VMMC |
| Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings | | | | | | | | | 10 |
| % of priority locations/districts with dedicated programs for AGYW & partners | id | Condom use with your require and trace (6/) | Women 15-24 | 46 | | Wo | men 15-24 | 0.8 | ART |
| Educational policies on HIV & sexuality education (secondary school) | Yes | Condom use with non-regular partners (%) | Men 15-24 | 78 | | | Men 15-24 | 0.6 | 10 |
| Laws requiring parental consent for adolescents to access HIV testing services | Yes, <12 | % who completed lower secondary education | Girls | 30 | | | Adults, 15+ | 2.7 | PrEP |
| Provider-initiated condom promotion integrated into SRH services | No/id | % of women experienced physical or sexual | Women 15-19 | id | | | | | 10 |
| HIV testing services integrated with SRH services | Yes | violence from husband/partner | Women 15-49 | 24 | | | | | |
| onyms: na not applicable; | | Very good | | | Low | | | | |
| IIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men; | | Good | | | Very low | | | | |
| 1 sexual and reproductive health. | | Medium | | id | Insufficient | data | | | |