Select country from dropdown menu							
South Africa	ZAF	Based on most recent available data.	Version	2.02.		2023	
utput (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		
Number of condoms distributed and sold / year (in millions)	404	Condom use with non-regular partners (%)	Women 15-49 Men 15-49	60	(ages 15+ trend vs. 2020 and 2025 targets)		2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	20			68	400'000		• 2020, 2022
Estimated condom distribution need met (%)	63				350'000 340 000		2020 target
Men and boys (including VMMC)					300'000		2025 target
Number of VMMCs performed / year (in thousands)	150	% of 2020 VMMC target achieved	Men	58	250'000		
% of annual VMMC target achieved	100	National male circumcision prevalence (%)	Men 15-24	60	200'000 180 0	00	2020 and 202
			Men 15-49	id	150'000	150'000	targets repres the country's
ARV-based prevention					100'000	69 000	required contribution 1
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	61	50'000 86 000	•	global targets
Number of people who received PrEP at least once in the past 12 months	406170		Women 15+	74	<u> </u>		75% reductio 2020 and 82.
% of PLHIV on ART	75	% of PLHIV virally surpressed	Men 15+	62	2010 2015 2020	2025	reduction by
Key populations					HIV	/ prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	146000	Condom use / last paid sex (%)	Sex workers	id	Sex workers <25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	34		Men 15-49	83	Sex workers all ages	45.0	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	70			
Gay men and other men who have sex with men (MSM)							Summai
Population size estimate for men who have sex with men	309700	Condom use / last anal sex (%)	MSM	72	MSM <25 years	id	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	10	% on ART	MSM LHIV	44	MSM all ages	42.9	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						6
People who inject drugs (PWID)							Sex work
Population size estimate for people who inject drugs	82500	% with safe injecting practices	514/15	id	PWID <25 years	id	6
% of PWID who received at least two HIV prevention interventions (past 3 mo)	17	% on opioid substitution therapy	PWID	1	PWID all ages	19.5	MSM
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	id			4
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	id			2
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	id			Condom
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id			6
Criminalization of transgender people	No		Transgender	id			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							8
% of priority locations/districts with dedicated programs for AGYW & partners	50		Women 15-24	61	Women 15-24	8.7	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	73	Men 15-24	3.6	8
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	91	Adults, 15+	17.3	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	30			6
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	30			
onyms: na not applicable;		Very good			Low		
IIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		
H sexual and reproductive health.		Medium		id	Insufficient data		