

Select country from dropdown menu

South Africa

ZAF

Based on most recent available data.

Version 2.02.


2024

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms			Condom use with non-regular partners (%)			New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)		
Number of condoms distributed and sold / year (in millions)	548		Women 15-49	60		2010	320 000	
Number of condoms distributed/sold per couple-year (age range 15-64)	27		Men 15-49	68		2015		
Estimated condom distribution need met (%)	78					2020	160 000	2020 target
Men and boys (including VMMC)			% of 2020 VMMC target achieved			2025 target		
Number of VMMCs performed / year (in thousands)	150		Men	58		2025	63 000	
% of annual VMMC target achieved	100		Men 15-24	60				
			Men 15-49	id				
ARV-based prevention			National male circumcision prevalence (%)			HIV prevalence		
Composite PrEP score (0-10)	10		All pop.	id				
Number of people who received PrEP at least once in the past 12 months	803171		Women 15+	75				
% of PLHIV on ART	77		Men 15+	64				
Key populations			Sex workers (SW)			Sex workers <25 years		
Population size estimate for sex workers	id		Sex workers	id		Sex workers all ages	45.0	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	34		Men 15-49	83				
Prevention strategy includes core elements of SW prevention package	> Half		% on ART	70				
Gay men and other men who have sex with men (MSM)			MSM <25 years			MSM all ages		
Population size estimate for men who have sex with men	309700		MSM	72		MSM <25 years	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	10		MSM LHIV	44		MSM all ages	42.9	
Prevention strategy includes core elements of MSM prevention package	> Half							
People who inject drugs (PWID)			PWID <25 years			PWID all ages		
Population size estimate for people who inject drugs	id		PWID	95		PWID <25 years	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	17		% with safe injecting practices	1		PWID all ages	19.5	
Prevention strategy includes core elements of PWID harm reduction package	< Half		% on opioid substitution therapy	id				
			% on ART	id				
Structural barriers and enablers			% of people who avoided health care because of stigma and discrimination			Summary Scores		
Criminalization of sex work	Yes		Sex workers	id		AGYW	6	
Criminalization of same-sex sexual acts	No		MSM	id		Sex workers	6	
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		MSM	4	
Criminalization of transgender people	No		Transgender	id		PWID	5	
						Condoms	7	
						VMMC	8	
						ART	8	
						PrEP	10	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings			Condom use with non-regular partners (%)			Women 15-24		
% of priority locations/districts with dedicated programs for AGYW & partners	52		Women 15-24	61		Women 15-24	8.4	
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	73		Men 15-24	3.8	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12		% who completed lower secondary education	91		Adults, 15+	16.6	
Provider-initiated condom promotion integrated into SRH services	Yes		% of women experienced physical or sexual violence from husband/partner	30				
HIV testing services integrated with SRH services	Yes		Women 15-19	30				
			Women 15-49	30				



2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

Acronyms: na ... not applicable;
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
 SRH ... sexual and reproductive health.

Very good
 Good
 Medium

Low
 Very low
 Insufficient data