South Sudan	SSD	Based on most recent available data.	Version	2.02.			2023	
Output (coverage)		Outcome (service use/behaviour)			Impact			
Condoms					New HIV infections			_
Number of condoms distributed and sold / year (in millions)	0		Women 15-49	id	(ages 15+ trend vs. 2020	and 2025 targ	jets)	2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	0	Condom use with non-regular partners (%)	Men 15-49	id	16'000			• 2020, 2022
Estimated condom distribution need met (%)	id				14'000 13 000			2020 target
Men and boys (including VMMC)					12'000			2025 target
Number of VMMCs performed / year (in thousands)	12	% of 2020 VMMC target achieved	Men	1	10'000	9 600	9'000	
% of annual VMMC target achieved	3	National male circumcision prevalence (%)	Men 15-24	id	8'000			2020 and 202 targets repre
			Men 15-49	id	6'000			the country's
ARV-based prevention					4'000		2 600	required contribution
Composite PrEP score (0-10)	3	% of national PrEP need met	All pop.	1	2'000	3 300	•	global target
Number of people who received PrEP at least once in the past 12 months	153		Women 15+	id	-			75% reductio 2020 and 82.
% of PLHIV on ART	32	% of PLHIV virally surpressed	Men 15+	id	2010 2015	2020	2025	reduction by
Key populations						HI	/ prevalence	2025 against 2010 as a
Sex workers (SW)								baseline.
Population size estimate for sex workers	45000	Condem use (last reid sey (9))	Sex workers	35	Sex workers	<25 years	3.9	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	39	Condom use / last paid sex (%)	Men 15-49	id	Sex worke	rs all ages	16.0	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	75				
Gay men and other men who have sex with men (MSM)								Summa
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	id	MSM	<25 years	id	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id	% on ART	MSM LHIV	id	MSI	M all ages	id	AGYW
Prevention strategy includes core elements of MSM prevention package	None							1
People who inject drugs (PWID)								Sex work
Population size estimate for people who inject drugs	id	% with safe injecting practices	DMUD	id	PWID	<25 years	id	2
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWI	D all ages	id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id				id
Structural barriers and enablers								PWID
Criminalization of sex work	Yes		Sex workers	12				id
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of	MSM	id				Condor
Criminalization of drug use/consumption or possession for personal use	id	stigma and discrimination	PWID	id				id
Criminalization of transgender people	No		Transgender	id				VMM
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								0
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom uses with non-require portners (%)	Women 15-24	id	Worr	nen 15-24	0.9	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	id	N	len 15-24	0.5	3
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <18	% who completed lower secondary education	Girls	10	Ad	dults, 15+	1.9	PrEP
Provider-initiated condom promotion integrated into SRH services	No/id	% of women experienced physical or sexual	Women 15-19	id				3
HIV testing services integrated with SRH services	Partial	violence from husband/partner	Women 15-49	id				
nyms: na not applicable;		Very good			Low			
people living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low			