Select country from dropdown menu							
United Republic of Tanzania	TZA	Based on most recent available data.	Version	2.02.		2023	
Output (coverage)		Outcome (service use/behaviour)			Impact		
Condoms					New HIV infections		
Number of condoms distributed and sold / year (in millions)	134	Condom use with non-regular partners (%)	Women 15-49	28	(ages 15+ trend vs. 2020 and 2025 targets)		<ul><li>2010 baseline</li><li>2020, 2022</li></ul>
Number of condoms distributed/sold per couple-year (age range 15-64)	8		Men 15-49	35	100'000		
Estimated condom distribution need met (%)	42				90'000 84 000		◆2020 target
Men and boys (including VMMC)					80′000		2025 target
Number of VMMCs performed / year (in thousands)	454	% of 2020 VMMC target achieved	Men	100	70'000 60'000		
% of annual VMMC target achieved	100		Men 15-24	82	50/000		2020 and 202
National male circum	National male circumcision prevalence (%)	Men 15-49	80	41 000		targets repres the country's	
ARV-based prevention						26′ <b>DD00</b> 00	required contribution t
Composite PrEP score (0-10)	10	% of national PrEP need met	All pop.	75	20'000 10'000 21 000	•	global targets
Number of people who received PrEP at least once in the past 12 months	162477	% of PLHIV virally surpressed	Women 15+	96	-		75% reductio 2020 and 82.
% of PLHIV on ART	94		Men 15+	90	2010 2015 2020	2025	reduction by
Key populations					HIV	prevalence	2025 against 2010 as a
Sex workers (SW)							baseline.
Population size estimate for sex workers	id	Condom use / last paid sex (%)	Sex workers	72	Sex workers <25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	90		Men 15-49	id	Sex workers all ages	15.4	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id			
Gay men and other men who have sex with men (MSM)							Summai
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	id	MSM <25 years	id	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	4	% on ART	MSM LHIV	id	MSM all ages	8.4	AGYW
Prevention strategy includes core elements of MSM prevention package	> Half						3
People who inject drugs (PWID)							Sex work
Population size estimate for people who inject drugs	36000	% with safe injecting practices	D) 4 // D	id	PWID <25 years	id	7
% of PWID who received at least two HIV prevention interventions (past 3 mo)	11	% on opioid substitution therapy	PWID	26	PWID all ages	id	MSM
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	id			id
Structural barriers and enablers							PWID
Criminalization of sex work	Yes		Sex workers	id			1
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of	MSM	id			Condon
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id			4
Criminalization of transgender people	No		Transgender	id			VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							10
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular next next (n/)	Women 15-24	id	Women 15-24	1.7	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	id	Men 15-24	0.9	9
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	27	Adults, 15+	4.7	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id			10
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id			
onyms: na not applicable;		Very good			Low		<del>_</del>
HVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low		
H sexual and reproductive health.		Medium		id	Insufficient data		