Select country from dropdown menu								
Uganda	UGA	Based on most recent available data.	Version	2.02.		2	023	
Output (coverage)		Outcome (service use/behaviour)			Impact			
Condoms					New HIV infe	2040		
Number of condoms distributed and sold / year (in millions)	131	Condom use with non-regular neutrors (0/)	Women 15-49 37		(ages 15+ trend vs. 2020 and 2025 ta			2010 baseline
Number of condoms distributed/sold per couple-year (age range 15-64)	11	Condom use with non-regular partners (%)	Men 15-49	58	70′000 _66 000			<b>2</b> 020, 2022
Estimated condom distribution need met (%)	60				60'000			◆2020 target
Men and boys (including VMMC)					50'000	46,000		◆ 2025 target
Number of VMMCs performed / year (in thousands)	517	% of 2020 VMMC target achieved	Men	24		46 000 46	′000	
% of annual VMMC target achieved	77	National male circumcision prevalence (%)	Men 15-24	49	40'000			2020 and 2025 targets represent
		National male circumcision prevalence (%)	Men 15-49	46	30'000			the country's
ARV-based prevention					20'000	•	13 000	required contribution to
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id	10'000	16 000	<b>*</b>	global targets, a
Number of people who received PrEP at least once in the past 12 months	195743	0/ -f DILIN/ 2 II	Women 15+	83				75% reduction by 2020 and 82.5%
% of PLHIV on ART	84	% of PLHIV virally surpressed	Men 15+	75	2010 2015	2020	2025	reduction by
Key populations						HIV pre	valence	2025 against 2010 as a
Sex workers (SW)								baseline.
Population size estimate for sex workers	130000	6 1 (1)	Sex workers	id	Sex workers <	25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	40	Condom use / last paid sex (%)	Men 15-49	73	Sex workers	all ages	id	
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	65				
Gay men and other men who have sex with men (MSM)								Summary
Population size estimate for men who have sex with men	24100	Condom use / last anal sex (%)	MSM	id	MSM <	25 years	id	Scores
% of MSM who received at least two HIV prevention interventions (past 3 mo)	20	% on ART	MSM LHIV	66	MSM	all ages 1	2.7	AGYW
Prevention strategy includes core elements of MSM prevention package	id					_		4
People who inject drugs (PWID)								Sex workers
Population size estimate for people who inject drugs	7400	% with safe injecting practices		id	PWID <	25 years	id	6
% of PWID who received at least two HIV prevention interventions (past 3 mo)	5	% on opioid substitution therapy	PWID	id	PWID	all ages	id	MSM
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	78				id
Structural barriers and enablers								PWID
Criminalization of sex work	Yes		Sex workers	9				0
Criminalization of same-sex sexual acts	Yes	% of people who avoided health care because of	MSM	id				Condoms
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id				5
Criminalization of transgender people	No		Transgender	id				VMMC
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings			Ü					5
% of priority locations/districts with dedicated programs for AGYW & partners	38		Women 15-24	43	Wome	en 15-24	2.5	ART
Educational policies on HIV & sexuality education (secondary school)	Yes	Condom use with non-regular partners (%)	Men 15-24	57			L.O	8
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	23			5.5	PrEP
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	Women 15-19	id				6
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	id				
Acronyms: na not applicable;		Very good			Low			
PLHIVpeople living with HIV; LHIVliving with HIV; YWyoung women; YMyoung men;		Good			Very low			
SRH sexual and reproductive health.		Medium		id	Insufficient data			<u> </u>