

Coverage

Outcome

Impact

Summary

Prevention strategy & systems

Comments

Scores

National and sub-national HIV prevention targets in place for all relevant pillars

Some

National standard operating procedures (detailed guidance) for all relevant pillars

Yes

Condoms

Number of condoms distributed and sold / year (in millions)

290

Number of condoms distributed and sold per man 15-64 /year

6

% of condom distribution need met

25

Voluntary medical male circumcision (VMMC)

Number of VMMCs performed / year (in thousands)

na

% of annual VMMC target achieved

na

ARV-based prevention

Composite PrEP score (0-10)

5

Number of people who received PrEP at least once in the past 12 months

360

Key populations

Sex workers (SW)

Population size estimate for sex workers (in thousands)

100

% of sex workers who received at least two HIV prevention interventions (past 3 mo)

44

% of sex workers who avoided health care because of stigma / discrimination

id

Men who have sex with men (MSM)

Population size estimate for men who have sex with men (in thousands)

26

% of MSM who received at least two HIV prevention interventions (past 3 mo)

25

% of MSM who avoided health care because of stigma / discrimination

id

People who inject drugs (PWID)

Population size estimate for people who inject drugs (in thousands)

45

% of PWID who received at least two HIV prevention interventions (past 3 mo)

15

% of PWID who avoided health care because of stigma / discrimination

id

Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings

% of priority districts (admin. areas) with dedicated programs for AGYW & partners

id

% of girls who completed lower secondary education

45

% of women 15-49 who experienced physical or sexual violence f.a husband/ partner

11

Condom use with non-regular partners (%)

Women 15-49

40

Men 15-49

58

% of 2021 VMMC target achieved

Men

na

National MC prevalence (%)

Men 15-24

99

Men 15-49

99

% of national PrEP need met

All pop.

id

% of PLHIV on ART

All PLHIV

33

% of PLHIV virally suppressed

All PLHIV

25

Condom use / last paid sex (%)

Sex workers

98

Condom use / last paid sex (%)

Men 15-49

66

Condom use / last anal sex (%)

MSM

51

% with safe injecting practices

PWID

71

% of PWID on OST

PWID

id

Condom use with non-regular partners (%)

YW 15-24

43

% who had multiple sexual partners

YM 15-24

58

% who had multiple sexual partners

AGYW 15-24*

2


% who had multiple sexual partners

Men 15-49**

13

* among sexually active AGYW; ** among all men

New HIV infections (ages 15+, in 1000s)



AGYW&MPs

4

SW

6

MSM

4

PWID

3

Condoms

4

VMMC

na

PrEP

5

HIV prevalence

Sex workers <25 years

id

Sex workers all ages

14

MSM <25 years

19

MSM all ages

23

PWID <25 years

2

PWID all ages

3

Young women 15-24

1,7

Young men 15-24

1,2

Adults 15-49

2,8

Acronyms

na ... not applicable

OST ... opioid substitution therapy

Very good

Good

Medium

Low

Very low

Insufficient data