

# Angola

2019

Based on most recent available data.

Version 2.01.

## Coverage

## Outcome

## Impact

## Summary

Condoms		Condom use with non-regular partners (%)	Women 15-49		New HIV infections (ages 15+, in 1000s)	Scores
Number of condoms distributed and sold / year (in millions)	54		Men 15-49	29		
Number of condoms distributed and sold per man 15-64 /year	8		Men 15-49	53		AGYW&MPs
% of condom distribution need met	20					3
Voluntary medical male circumcision (VMMC)		% of 2020 VMMC target achieved	Men		2018; 21	SW
Number of VMMCs performed / year (in thousands)	na		Men	na		5
% of annual VMMC target achieved	na	National MC prevalence (%)	Men 15-24	96		MSM
			Men 15-49	96		id
ARV-based prevention		% of national PrEP need met	All pop.	id		PWID
Composite PrEP score (0-10)	0		% of PLHIV on ART	All PLHIV	27	
Number of people who received PrEP at least once in the past 12 months	0	% of PLHIV virally suppressed	All PLHIV	15		Condoms
						3
						VMMC
						na
						PrEP
						0
Key populations		Condom use / last paid sex (%)	Sex workers (SW)		HIV prevalence	
Population size estimate for sex workers (in thousands)	125		Sex workers	72	Sex workers <25 years	7
% of SWs who received at least two HIV prevention interventions (past 3 mo)	43	Men 15-49	71	Sex workers all ages	8	
Men who have sex with men (MSM)		Condom use / last anal sex (%)	MSM	59	MSM <25 years	8
Population size estimate for men who have sex with men (in thousands)	106				MSM all ages	2
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id					
People who inject drugs (PWID)		% with safe injecting practices	PWID	id	PWID <25 years	id
Population size estimate for people who inject drugs (in thousands)	id	% of PWID on OST	PWID	id	PWID all ages	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id					
Structural barriers and enablers						
Training on HIV-related rights & non-discrimination: law enforcement officials	Sub-national	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id		
			PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings		Condom use with non-regular partners (%)	YW 15-24	33	Young women 15-24	1.2
% of priority locations/districts with dedicated programs for AGYW & partners	id		YM 15-24	52	Young men 15-24	0.3
% of girls who completed lower secondary education	id				Adults 15-49	2.0
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	0					
Background population-level factors		% who had multiple sexual partners	Men 15-49	18		
% of women 15-49 experienced physical or sexual violence f. husband/partner	26					
% of married women who decide on their own health care	22					
Acronyms		Very good			Low	
na ... not applicable		Good			Very low	
OST ... opioid substitution therapy		Medium		id	Insufficient data	