

Cote d'Ivoire

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage				Outcome				Impact		Summary																			
Condoms				Condom use with non-regular partners (%)	Women 15-49	31	New HIV infections (ages 15+, in 1000s)	2010	2015	2018; 17	2020	6	<table border="1"> <thead> <tr> <th colspan="2">Scores</th> </tr> </thead> <tbody> <tr> <td>AGYW&MPs</td> <td>7</td> </tr> <tr> <td>SW</td> <td>8</td> </tr> <tr> <td>MSM</td> <td>6</td> </tr> <tr> <td>PWID</td> <td>id</td> </tr> <tr> <td>Condoms</td> <td>3</td> </tr> <tr> <td>VMMC</td> <td>na</td> </tr> <tr> <td>PrEP</td> <td>5</td> </tr> </tbody> </table>	Scores		AGYW&MPs	7	SW	8	MSM	6	PWID	id	Condoms	3	VMMC	na	PrEP	5
Scores																													
AGYW&MPs	7																												
SW	8																												
MSM	6																												
PWID	id																												
Condoms	3																												
VMMC	na																												
PrEP	5																												
Number of condoms distributed and sold / year (in millions)	34	Men 15-49	42																										
Number of condoms distributed and sold per man 15-64 / year	5																												
% of condom distribution need met	17																												
Voluntary medical male circumcision (VMMC)				% of 2020 VMMC target achieved	Men	na																							
Number of VMMCs performed / year (in thousands)	na	Men 15-24	96																										
% of annual VMMC target achieved	na	Men 15-49	97																										
ARV-based prevention				% of national PrEP need met	All pop.	id																							
Composite PrEP score (0-10)	5	% of PLHIV on ART	85																										
Number of people who received PrEP at least once in the past 12 months	100	% of PLHIV virally suppressed	76																										
Key populations				reported by				HIV prevalence																					
<i>Sex workers (SW)</i>					Sex workers	94	Sex workers <25 years	2																					
Population size estimate for sex workers (in thousands)	51	Condom use / last paid sex (%)	94	Men 15-49	42	Sex workers all ages	11																						
% of SWs who received at least two HIV prevention interventions (past 3 mo)	96	Condom use / last paid sex (%)	42																										
<i>Men who have sex with men (MSM)</i>					MSM	75	MSM <25 years	10																					
Population size estimate for men who have sex with men (in thousands)	30	Condom use / last anal sex (%)	75			MSM all ages	12																						
% of MSM who received at least two HIV prevention interventions (past 3 mo)	39																												
<i>People who inject drugs (PWID)</i>																													
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	id			PWID <25 years	id																						
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	id			PWID all ages	5																						
<i>Structural barriers and enablers</i>																													
Training on HIV-related rights & non-discrimination: law enforcement officials	id	% of people who avoided health care because of stigma and discrimination	Sex workers	6																									
Training on HIV-related rights & non-discrimination: health care workers	id		MSM	22																									
			PWID	id																									
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings				Condom use with non-regular partners (%)	YW 15-24	48	Young women 15-24	0.4																					
% of priority locations/districts with dedicated programs for AGYW & partners	100	YM 15-24	67		Young men 15-24	0.6																							
% of girls who completed lower secondary education	28				Adults 15-49	2.6																							
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	id																												
<i>Background population-level factors</i>																													
% of women 15-49 experienced physical or sexual violence f. husband/partner	31	% who had multiple sexual partners	24																										
% of married women who decide on their own health care	10																												
Acronyms		Very good				Low																							
na ... not applicable		Good				Very low																							
OST ... opioid substitution therapy		Medium			id	Insufficient data																							

