

Cameroon

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage				Outcome				Impact		Summary																				
Condoms				Condom use with non-regular partners (%)	Women 15-49	55	New HIV infections (ages 15+, in 1000s)	28	2010	2015	2018; 19	2020	7	<table border="1"> <thead> <tr> <th colspan="2">Scores</th> </tr> </thead> <tbody> <tr> <td>AGYW&MPs</td> <td>6</td> </tr> <tr> <td>SW</td> <td>5</td> </tr> <tr> <td>MSM</td> <td>5</td> </tr> <tr> <td>PWID</td> <td>id</td> </tr> <tr> <td>Condoms</td> <td>4</td> </tr> <tr> <td>VMMC</td> <td>na</td> </tr> <tr> <td>PrEP</td> <td>2</td> </tr> </tbody> </table>	Scores		AGYW&MPs	6	SW	5	MSM	5	PWID	id	Condoms	4	VMMC	na	PrEP	2
Scores																														
AGYW&MPs	6																													
SW	5																													
MSM	5																													
PWID	id																													
Condoms	4																													
VMMC	na																													
PrEP	2																													
Number of condoms distributed and sold / year (in millions)	69	Men 15-49	74																											
Number of condoms distributed and sold per man 15-64 / year	10																													
% of condom distribution need met	26																													
Voluntary medical male circumcision (VMMC)				% of 2020 VMMC target achieved	Men	na																								
Number of VMMCs performed / year (in thousands)	na	Men 15-24	93																											
% of annual VMMC target achieved	na	Men 15-49	94																											
ARV-based prevention				% of national PrEP need met	All pop.	id																								
Composite PrEP score (0-10)	2																													
Number of people who received PrEP at least once in the past 12 months	0	% of PLHIV on ART	53																											
		% of PLHIV virally suppressed	33																											
Key populations				reported by				HIV prevalence																						
<i>Sex workers (SW)</i>					Sex workers	96	Sex workers <25 years	28																						
Population size estimate for sex workers (in thousands)	71	Condom use / last paid sex (%)		Men 15-49	81	Sex workers all ages	24																							
% of SWs who received at least two HIV prevention interventions (past 3 mo)	23	Condom use / last paid sex (%)																												
<i>Men who have sex with men (MSM)</i>					MSM	78	MSM <25 years	29																						
Population size estimate for men who have sex with men (in thousands)	7	Condom use / last anal sex (%)				MSM all ages	21																							
% of MSM who received at least two HIV prevention interventions (past 3 mo)	17																													
<i>People who inject drugs (PWID)</i>							PWID <25 years	id																						
Population size estimate for people who inject drugs (in thousands)	2	% with safe injecting practices		PWID	id	PWID all ages	id																							
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST		PWID	id																									
<i>Structural barriers and enablers</i>																														
Training on HIV-related rights & non-discrimination: law enforcement officials	Limited	% of people who avoided health care because of stigma and discrimination		Sex workers	5																									
Training on HIV-related rights & non-discrimination: health care workers	Limited		MSM	14																										
			PWID	id																										
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings				Condom use with non-regular partners (%)	YW 15-24	61	Young women 15-24	1.8																						
% of priority locations/districts with dedicated programs for AGYW & partners	id	YM 15-24	77		Young men 15-24	0.7																								
% of girls who completed lower secondary education	68				Adults 15-49	3.6																								
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	63																													
<i>Background population-level factors</i>																														
% of women 15-49 experienced physical or sexual violence f. husband/partner	id	% who had multiple sexual partners		Men 15-49	29																									
% of married women who decide on their own health care	16																													
Acronyms				Very good		Low																								
na ... not applicable		Good				Very low																								
OST ... opioid substitution therapy		Medium		id		Insufficient data																								