

Eswatini

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary															
Condoms																					
Number of condoms distributed and sold / year (in millions)	26	Condom use with non-regular partners (%)	Women 15-49	54	New HIV infections (ages 15+, in 1000s)	<table border="1"> <thead> <tr> <th>Scores</th> </tr> </thead> <tbody> <tr> <td>AGYW&MPs</td> </tr> <tr> <td>7</td> </tr> <tr> <td>SW</td> </tr> <tr> <td>id</td> </tr> <tr> <td>MSM</td> </tr> <tr> <td>4</td> </tr> <tr> <td>PWID</td> </tr> <tr> <td>3</td> </tr> <tr> <td>Condoms</td> </tr> <tr> <td>8</td> </tr> <tr> <td>VMMC</td> </tr> <tr> <td>4</td> </tr> <tr> <td>PrEP</td> </tr> <tr> <td>9</td> </tr> </tbody> </table>	Scores	AGYW&MPs	7	SW	id	MSM	4	PWID	3	Condoms	8	VMMC	4	PrEP	9
Scores																					
AGYW&MPs																					
7																					
SW																					
id																					
MSM																					
4																					
PWID																					
3																					
Condoms																					
8																					
VMMC																					
4																					
PrEP																					
9																					
Number of condoms distributed and sold per man 15-64 / year	82	Men 15-49	67																		
% of condom distribution need met	100																				
Voluntary medical male circumcision (VMMC)																					
Number of VMMCs performed / year (in thousands)	14	% of 2020 VMMC target achieved	Men	26																	
% of annual VMMC target achieved	40	National MC prevalence (%)	Men 15-24	12																	
			Men 15-49	10																	
ARV-based prevention																					
Composite PrEP score (0-10)	9	% of national PrEP need met	All pop.	id																	
Number of people who received PrEP at least once in the past 12 months	2176	% of PLHIV on ART	All PLHIV	86																	
		% of PLHIV virally suppressed	All PLHIV	81																	
Key populations																					
<i>Sex workers (SW)</i>			reported by		HIV prevalence																
Population size estimate for sex workers (in thousands)	12	Condom use / last paid sex (%)	Sex workers	83	Sex workers <25 years	64															
% of SWs who received at least two HIV prevention interventions (past 3 mo)	id	Condom use / last paid sex (%)	Men 15-49	id	Sex workers all ages	61															
<i>Men who have sex with men (MSM)</i>																					
Population size estimate for men who have sex with men (in thousands)	6	Condom use / last anal sex (%)	MSM	58	MSM <25 years	id															
% of MSM who received at least two HIV prevention interventions (past 3 mo)	24				MSM all ages	13															
<i>People who inject drugs (PWID)</i>																					
Population size estimate for people who inject drugs (in thousands)	0	% with safe injecting practices	PWID	id	PWID <25 years	id															
% of PWID who received at least two HIV prevention interventions (past 3 mo)	31	% of PWID on OST	PWID	id	PWID all ages	id															
<i>Structural barriers and enablers</i>																					
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	21																	
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	35																	
			PWID	id																	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings																					
% of priority locations/districts with dedicated programs for AGYW & partners	71	Condom use with non-regular partners (%)	YW 15-24	71	Young women 15-24	15.9															
% of girls who completed lower secondary education	52		YM 15-24	93	Young men 15-24	3.1															
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	100				Adults 15-49	27.3															
<i>Background population-level factors</i>																					
% of women 15-49 experienced physical or sexual violence f. husband/partner	id	% who had multiple sexual partners	Men 15-49	14																	
% of married women who decide on their own health care	36																				
Acronyms																					
na ... not applicable		Very good			Low																
OST ... opioid substitution therapy		Good			Very low																
		Medium		id	Insufficient data																

