

Ethiopia

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary	
Condoms							
Number of condoms distributed and sold / year (in millions)	173	Condom use with non-regular partners (%)	Women 15-49	20	New HIV infections (ages 15+, in 1000s)	Scores	
Number of condoms distributed and sold per man 15-64 / year	6		Men 15-49	51		AGYW&MPs	2
% of condom distribution need met	62					SW	id
Voluntary medical male circumcision (VMMC)							
Number of VMMCs performed / year (in thousands)	23	% of 2020 VMMC target achieved	Men	100		MSM	id
% of annual VMMC target achieved	100	National MC prevalence (%)	Men 15-24	88		PWID	id
			Men 15-49	91		Condoms	5
ARV-based prevention							
Composite PrEP score (0-10)	0	% of national PrEP need met	All pop.	id	VMMC	10	
Number of people who received PrEP at least once in the past 12 months	0	% of PLHIV on ART	All PLHIV	65	PrEP	0	
		% of PLHIV virally suppressed	All PLHIV	70			
Key populations							
<i>Sex workers (SW)</i>				reported by		HIV prevalence	
Population size estimate for sex workers (in thousands)	160	Condom use / last paid sex (%)	Sex workers	98	Sex workers <25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	id	Condom use / last paid sex (%)	Men 15-49	81	Sex workers all ages	24	
<i>Men who have sex with men (MSM)</i>							
Population size estimate for men who have sex with men (in thousands)	id	Condom use / last anal sex (%)	MSM	id	MSM <25 years	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id				MSM all ages	id	
<i>People who inject drugs (PWID)</i>							
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	PWID	id	PWID all ages	id	
<i>Structural barriers and enablers</i>							
Training on HIV-related rights & non-discrimination: law enforcement officials	Limited	% of people who avoided health care because of stigma and discrimination	Sex workers	id			
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id			
			PWID	id			
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	YW 15-24	24	Young women 15-24	0.4	
% of girls who completed lower secondary education	17		YM 15-24	55	Young men 15-24	0.2	
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	72				Adults 15-49	1.0	
<i>Background population-level factors</i>							
% of women 15-49 experienced physical or sexual violence f. husband/partner	20	% who had multiple sexual partners	Men 15-49	3			
% of married women who decide on their own health care	15						
Acronyms		Very good			Low		
na ... not applicable		Good			Very low		
OST ... opioid substitution therapy		Medium		id	Insufficient data		