

Ghana

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Condoms				Condom use with non-regular partners (%)		New HIV infections (ages 15+, in 1000s)		Scores	
Number of condoms distributed and sold / year (in millions)	37	Women 15-49	17	2010	17	AGYW&MPs	3		
Number of condoms distributed and sold per man 15-64 / year	4	Men 15-49	39	2015	17	SW	6		
% of condom distribution need met	17			2018	17	MSM	6		
Voluntary medical male circumcision (VMMC)				National MC prevalence (%)					
Number of VMMCs performed / year (in thousands)	na	Men	na			PWID	4		
% of annual VMMC target achieved	na	Men 15-24	96			id			
		Men 15-49	96			Condoms	2		
ARV-based prevention				% of PLHIV virally suppressed					
Composite PrEP score (0-10)	0	All pop.	id			VMMC	na		
Number of people who received PrEP at least once in the past 12 months	0	All PLHIV	34			PrEP	0		
Key populations				HIV prevalence					
Sex workers (SW)				reported by					
Population size estimate for sex workers (in thousands)	65	Sex workers	90	Sex workers <25 years	3				
% of SWs who received at least two HIV prevention interventions (past 3 mo)	48	Men 15-49	44	Sex workers all ages	7				
Men who have sex with men (MSM)									
Population size estimate for men who have sex with men (in thousands)	55	MSM	60	MSM <25 years	id				
% of MSM who received at least two HIV prevention interventions (past 3 mo)	55			MSM all ages	18				
People who inject drugs (PWID)									
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	id	PWID <25 years	id				
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	id	PWID all ages	id				
Structural barriers and enablers									
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	id					
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id					
			PWID	id					
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings									
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	YW 15-24	19	Young women 15-24	1.0			
% of girls who completed lower secondary education	50		YM 15-24	39	Young men 15-24	0.4			
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	27				Adults 15-49	1.7			
Background population-level factors									
% of women 15-49 experienced physical or sexual violence f. husband/partner	id	% who had multiple sexual partners	Men 15-49	14					
% of married women who decide on their own health care	27								
Acronyms		Very good		Low					
na ... not applicable		Good		Very low					
OST ... opioid substitution therapy		Medium	id	Insufficient data					