

HIV prevention among key populations - thematic

2019

Version 2.01

Back
Stai

Thematic area	Indicator	Source	Country																								Scale										
			Angola	Botswana	Cameroon	Cote d'Ivoire	DR Congo	Ethiopia	Ghana	Kenya	Lesotho	Malawi	Mozambique	Namibia	Nigeria	South Africa	eSwatini	Tanzania	Uganda	Zambia	Zimbabwe	Brazil	China	India	Indonesia	Iran	Mexico	Myanmar	Pakistan	Ukraine							
Impact	HIV prevalence (sex workers, <25 years)	GAM 3.3A/ IBBS	7	id	28	2	5	id	3	id	id	id	id	10	id	64	id	id	id	21	2	0	1	4	1	5	4	2	1								
	HIV prevalence (sex workers, all ages)	GAM 3.3A/ IBBS	8	42	24	11	6	24	7	29	72	55	id	41	14	58	61	15	31	49	41	5	0	2	4	2	1	6	2	6							
	HIV prevalence (men who have sex with men, <25 years)	GAM 3.3B/ IBBS	8	id	29	10	2	id	id	12	id	id	id	id	19	id	id	15	id	id	id	10	6	3	24	id	13	4	4	7							
	HIV prevalence (men who have sex with men, all ages)	GAM 3.3B/ IBBS	2	15	21	12	3	id	18	18	33	7	id	id	23	18	13	8	13	id	31	18	7	3	26	id	13	6	5	8							
	HIV prevalence (people who inject drugs, <25 years)	GAM 3.3C/ IBBS	id	id	id	id	6	id	id	id	id	id	id	id	2	12	id	12	17	id	id	na	4	8	3	2	4	17	18	5							
	HIV prevalence (people who inject drugs, all ages)	GAM 3.3C/ IBBS	id	id	id	5	6	id	id	18	id	id	id	id	3	22	id	16	17	id	id	na	6	6	29	9	4	19	38	24							
	HIV prevalence (transgender people, <25 years)	GAM 3.3D/ IBBS	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	19	id	6	19	0	id	id	6	id							
HIV prevalence (transgender people, all ages)	GAM 3.3D/ IBBS	id	16	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	30	id	3	25	2	9	id	7	id								
Outcome	Condom use of sex workers with most recent client (reported by sex workers)	GAM 3.6A/ IBBS	72	76	96	94	69	98	90	92	62	65	id	id	98	86	83	70	69	79	96	86	94	91	68	59	87	81	51	94	95	90	80	70	0		
	Condom use at last anal sex among MSM	GAM 3.6B/ IBBS	59	78	78	75	77	id	60	55	46	44	id	id	51	98	58	14	39	id	id	64	85	84	81	id	73	77	26	78	90	80	70	60	0		
	Condom use among transgender people	GAM 3.6D/ IBBS	id	60	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	71	id	80	id	43	75	id	28	id	90	80	70	60	0		
	Condom use at last paid sex act (reported by men)	DHS or other survey	71	id	81	42	34	81	44	74	90	75	31	67	66	83	id	id	73	60	90	id	id	48	33	id	id	77	id	id	90	80	70	60	0		
	Safe injecting practices (people who inject drugs)	IBBS/ GAM 3.8	id	id	id	id	22	id	id	88	id	id	id	id	41	id	id	84	id	id	na	87	86	89	82	71	91	58	97	95	90	80	70	0			
	% of opioid users who receive opioid substitution therapy	GAM	id	id	id	id	id	id	id	26	id	id	id	id	id	id	id	21	id	id	id	na	id	20	11	12	id	17	id	4	50	40	25	10	0		
Output	Sex workers																																				
	Population size estimate for female sex workers in 1000s	GAM 3.2A/ Pop. size study	125	7	71	51	350	160	65	170	8	36	93	8	408	112	12	160	130	75	45	547	id	660	230	90	240	66	230	87							
	% of all sex workers who received at least two HIV prevention interventions in the past three months	GAM 3.7A/ IBBS/prg. data	43	60	23	96	id	id	48	51	55	68	51	56	44	45	id	id	id	47	44	22	id	73	40	29	id	id	1	48	90	75	60	50	0		
	% of sex workers who avoided health care because of stigma and discrimination	GAM	id	id	5	6	id	id	id	id	8	id	id	id	id	id	21	id	id	id	39	12	id	id	id	id	id	4	id	id	5	10	20	30	100		
	Men who have sex with men																																				
	Population size estimate for men who have sex with men in 1000s	GAM 3.2B/ Pop. size study/	106	3	7	30	id	id	55	33	6	43	41	7	227	300	6	50	22	67	id	2000	id	240	750	360	###	250	830	180							
	% of all MSM who who received at least two HIV prevention interventions in the past three months	GAM 3.7B/ IBBS/prg. data	id	64	17	39	id	id	55	id	40	65	4	44	25	33	24	28	id	13	id	30	id	59	20	id	id	id	1	24	90	75	60	50	0		
	% of men who have sex with men who avoided health care because of stigma and discrimination	GAM	id	id	14	22	id	id	id	id	8	id	id	id	id	id	35	id	id	id	id	63	id	id	id	id	id	8	id	id	5	10	20	30	100		
	People who inject drugs																																				
	Population size estimate for people who inject drugs available	GAM 3.2C/ Pop. size study/	id	id	2	id	id	id	id	16	id	id	14	id	170	75	0	30	7	27	id	na	id	130	34	200	110	93	110	350							
	% of all people who inject drugs who received at least two HIV prevention interventions in the past three months	GAM 3.7C/ IBBS/prg. data	id	id	id	id	id	id	id	34	id	id	15	id	id	24	31	21	8	id	id	na	id	76	44	35	id	34	2	65	90	75	60	50	0		
	Needles and syringes per person who inject drugs	GAM 3.9/ Prg. records	id	id	id	id	id	id	id	137	id	id	id	id	id	id	id	15	id	id	id	na	240	366	3	48	6	351	46	57	200	160	120	80	0		
% of people who inject drugs who avoided health care because of stigma and discrimination	GAM	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	64	id	id	na	id	id	id	id	id	id	id	id	id	5	10	20	30	100		
Transgender people																																					
Population size estimate for transgender people	GAM 3.2D/ Pop. size study/	3	0	id	id	id	id	id	4	id	id	id	id	id	id	id	id	id	id	id	id	id	70	id	id	id	id	id	52	id							
% of all TG people who who received at least two HIV prevention interventions in the past three months	GAM 3.7D/ IBBS/prg. data	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	59	id	id	id	id	id	id	id	4	id	90	75	60	50	0		
% of transgender people who avoided health care because of stigma and discrimination	GAM	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	5	10	20	30	100		

Legend: Very good (green), Good (light green), Medium (yellow), Low (orange), Very low (red), id (pink), Insufficient data (grey), na (grey), Not applicable (grey)