

Kenya

2019 Based on most recent available data.

Version 2.01.


Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary
Condoms						
Number of condoms distributed and sold / year (in millions)	182	Condom use with non-regular partners (%)	Women 15-49	57	New HIV infections (ages 15+, in 1000s)	Scores
Number of condoms distributed and sold per man 15-64 / year	13		Men 15-49	76		
% of condom distribution need met	52					AGYW&MPs
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	287	% of 2020 VMMC target achieved	Men	100		7
% of annual VMMC target achieved	100	National MC prevalence (%)	Men 15-24	91		SW
				Men 15-49	93	
ARV-based prevention						
Composite PrEP score (0-10)	10	% of national PrEP need met	All pop.	id		MSM
Number of people who received PrEP at least once in the past 12 months	30000	% of PLHIV on ART	All PLHIV	68		id
		% of PLHIV virally suppressed	All PLHIV	63		PWID
Key populations						
<i>Sex workers (SW)</i>			reported by		HIV prevalence	
Population size estimate for sex workers (in thousands)	170	Condom use / last paid sex (%)	Sex workers	92	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	51	Condom use / last paid sex (%)	Men 15-49	74	Sex workers all ages	29
<i>Men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	33	Condom use / last anal sex (%)	MSM	55	MSM <25 years	12
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id				MSM all ages	18
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	16	% with safe injecting practices	PWID	88	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	34	% of PWID on OST	PWID	26	PWID all ages	18
<i>Structural barriers and enablers</i>						
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id		
			PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	83	Condom use with non-regular partners (%)	YW 15-24	60	Young women 15-24	2.5
% of girls who completed lower secondary education	69		YM 15-24	77	Young men 15-24	1.5
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	45				Adults 15-49	4.7
<i>Background population-level factors</i>						
% of women 15-49 experienced physical or sexual violence f. husband/partner	25	% who had multiple sexual partners	Men 15-49	13		
% of married women who decide on their own health care	39					
Acronyms						
na ... not applicable		Very good			Low	
OST ... opioid substitution therapy		Good			Very low	
		Medium		id	Insufficient data	



Category	Score
AGYW&MPs	7
SW	7
MSM	id
PWID	6
Condoms	6
VMMC	10
PrEP	10