

Lesotho

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary	
Condoms							
Number of condoms distributed and sold / year (in millions)	30	Condom use with non-regular partners (%)	Women 15-49	66	New HIV infections (ages 15+, in 1000s)		Scores AGYW&MPs 8 SW 6 MSM 4 PWID id Condoms 8 VMMC 7 PrEP 9
Number of condoms distributed and sold per man 15-64 / year	45		Men 15-49	66			
% of condom distribution need met	88						
Voluntary medical male circumcision (VMMC)							
Number of VMMCs performed / year (in thousands)	26	% of 2020 VMMC target achieved	Men	41			
% of annual VMMC target achieved	68	National MC prevalence (%)	Men 15-24	70			
			Men 15-49	72			
ARV-based prevention							
Composite PrEP score (0-10)	9	% of national PrEP need met	All pop.	id			
Number of people who received PrEP at least once in the past 12 months	7300	% of PLHIV on ART	All PLHIV	61			
		% of PLHIV virally suppressed	All PLHIV	57			
Key populations							
Sex workers (SW)				reported by		HIV prevalence	
Population size estimate for sex workers (in thousands)	8	Condom use / last paid sex (%)	Sex workers	62	Sex workers <25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	55	Condom use / last paid sex (%)	Men 15-49	90	Sex workers all ages	72	
Men who have sex with men (MSM)							
Population size estimate for men who have sex with men (in thousands)	6	Condom use / last anal sex (%)	MSM	46	MSM <25 years	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	40				MSM all ages	33	
People who inject drugs (PWID)							
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	PWID	id	PWID all ages	id	
Structural barriers and enablers							
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	8			
Training on HIV-related rights & non-discrimination: health care workers	Sub-national		MSM	8			
			PWID	id			
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							
% of priority locations/districts with dedicated programs for AGYW & partners	100	Condom use with non-regular partners (%)	YW 15-24	82	Young women 15-24	10.8	
% of girls who completed lower secondary education	46		YM 15-24	79	Young men 15-24	4.1	
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	36				Adults 15-49	23.6	
Background population-level factors							
% of women 15-49 experienced physical or sexual violence f. husband/partner	1	% who had multiple sexual partners	Men 15-49	27			
% of married women who decide on their own health care	40						
Acronyms		Very good			Low		
na ... not applicable		Good			Very low		
OST ... opioid substitution therapy		Medium		id	Insufficient data		