

Namibia

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary		
Condoms								
Number of condoms distributed and sold / year (in millions)	29	Condom use with non-regular partners (%)	Women 15-49	66		Scores		
Number of condoms distributed and sold per man 15-64 /year	37		Men 15-49	80		AGYW&MPs	4	SW
% of condom distribution need met	88					6	MSM	id
Voluntary medical male circumcision (VMMC)								
Number of VMMCs performed / year (in thousands)	35	% of 2020 VMMC target achieved	Men	23	2	PWID	id	
% of annual VMMC target achieved	47	National MC prevalence (%)	Men 15-24	22	8	Condoms	8	
ARV-based prevention								
				Men 15-49	26	4	VMMC	4
Composite PrEP score (0-10)	10	% of national PrEP need met	All pop.	id	10	PrEP	10	
Number of people who received PrEP at least once in the past 12 months	5600	% of PLHIV on ART	All PLHIV	92				
		% of PLHIV virally suppressed	All PLHIV	87				
Key populations								
<i>Sex workers (SW)</i>				reported by		HIV prevalence		
Population size estimate for sex workers (in thousands)	8	Condom use / last paid sex (%)	Sex workers	id	Sex workers <25 years	id		
% of SWs who received at least two HIV prevention interventions (past 3 mo)	56	Condom use / last paid sex (%)	Men 15-49	67	Sex workers all ages	41		
<i>Men who have sex with men (MSM)</i>								
Population size estimate for men who have sex with men (in thousands)	7	Condom use / last anal sex (%)	MSM	id	MSM <25 years	id		
% of MSM who received at least two HIV prevention interventions (past 3 mo)	44				MSM all ages	id		
<i>People who inject drugs (PWID)</i>								
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id		
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	PWID	id	PWID all ages	id		
<i>Structural barriers and enablers</i>								
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	id				
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id				
			PWID	id				
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								
% of priority locations/districts with dedicated programs for AGYW & partners	12	Condom use with non-regular partners (%)	YW 15-24	68	Young women 15-24	4.5		
% of girls who completed lower secondary education	62		YM 15-24	82	Young men 15-24	2.6		
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	45				Adults 15-49	11.8		
<i>Background population-level factors</i>								
% of women 15-49 experienced physical or sexual violence f. husband/partner	20	% who had multiple sexual partners	Men 15-49	10				
% of married women who decide on their own health care	46							
Acronyms								
na ... not applicable		Very good			Low			
OST ... opioid substitution therapy		Good			Very low			
		Medium		id	Insufficient data			