

South Africa

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary
Condoms						
Number of condoms distributed and sold / year (in millions)	745	Condom use with non-regular partners (%)	Women 15-49	61		Scores
Number of condoms distributed and sold per man 15-64 / year	40		Men 15-49	69		AGYW&MPs
% of condom distribution need met	89					5
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	572	% of 2020 VMMC target achieved	Men	38		SW
% of annual VMMC target achieved	73	National MC prevalence (%)	Men 15-24	70		6
			Men 15-49	57		MSM
						7
ARV-based prevention						
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id		PWID
Number of people who received PrEP at least once in the past 12 months	8100	% of PLHIV on ART	All PLHIV	62		2
		% of PLHIV virally suppressed	All PLHIV	36		Condoms
						8
						VMMC
						7
						PrEP
						6
Key populations						
<i>Sex workers (SW)</i>			reported by		HIV prevalence	
Population size estimate for sex workers (in thousands)	112	Condom use / last paid sex (%)	Sex workers	86	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	45	Condom use / last paid sex (%)	Men 15-49	83	Sex workers all ages	58
<i>Men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	300	Condom use / last anal sex (%)	MSM	98	MSM <25 years	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	33				MSM all ages	18
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	75	% with safe injecting practices	PWID	id	PWID <25 years	12
% of PWID who received at least two HIV prevention interventions (past 3 mo)	24	% of PWID on OST	PWID	id	PWID all ages	22
<i>Structural barriers and enablers</i>						
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id		
			PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	29	Condom use with non-regular partners (%)	YW 15-24	47	Young women 15-24	10.2
% of girls who completed lower secondary education	86		YM 15-24	68	Young men 15-24	3.9
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	45				Adults 15-49	18.8
<i>Background population-level factors</i>						
% of women 15-49 experienced physical or sexual violence f. husband/partner	21	% who had multiple sexual partners	Men 15-49	18		
% of married women who decide on their own health care	41					
Acronyms						
na ... not applicable		Very good			Low	
OST ... opioid substitution therapy		Good			Very low	
		Medium		id	Insufficient data	