

Tanzania

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary
Condoms						
Number of condoms distributed and sold / year (in millions)	66	Condom use with non-regular partners (%)	Women 15-49	51	New HIV infections (ages 15+, in 1000s)	Scores
Number of condoms distributed and sold per man 15-64 / year	4		Men 15-49	60		
% of condom distribution need met	15					
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	886	% of 2020 VMMC target achieved	Men	84		AGYW&MPs
% of annual VMMC target achieved	100	National MC prevalence (%)	Men 15-24	82		3
			Men 15-49	80		SW
ARV-based prevention						
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id		id
Number of people who received PrEP at least once in the past 12 months	8000	% of PLHIV on ART	All PLHIV	71		MSM
		% of PLHIV virally suppressed	All PLHIV	62		2
Key populations						
<i>Sex workers (SW)</i>			reported by		HIV prevalence	
Population size estimate for sex workers (in thousands)	160	Condom use / last paid sex (%)	Sex workers	70	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	id	Condom use / last paid sex (%)	Men 15-49	id	Sex workers all ages	15
<i>Men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	50	Condom use / last anal sex (%)	MSM	14	MSM <25 years	15
% of MSM who received at least two HIV prevention interventions (past 3 mo)	28				MSM all ages	8
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	30	% with safe injecting practices	PWID	84	PWID <25 years	12
% of PWID who received at least two HIV prevention interventions (past 3 mo)	21	% of PWID on OST	PWID	21	PWID all ages	16
<i>Structural barriers and enablers</i>						
Training on HIV-related rights & non-discrimination: law enforcement officials	National	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Training on HIV-related rights & non-discrimination: health care workers	Limited		MSM	id		
			PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	19	Condom use with non-regular partners (%)	YW 15-24	49	Young women 15-24	2.2
% of girls who completed lower secondary education	27		YM 15-24	86	Young men 15-24	1.2
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	id				Adults 15-49	4.6
<i>Background population-level factors</i>						
% of women 15-49 experienced physical or sexual violence f. husband/partner	30	% who had multiple sexual partners	Men 15-49	21		
% of married women who decide on their own health care	16					
Acronyms						
na ... not applicable		Very good			Low	
OST ... opioid substitution therapy		Good			Very low	
		Medium		id	Insufficient data	