

Zimbabwe

2019 Based on most recent available data.

Version 2.01.

Coverage

Outcome

Impact

Summary

Coverage		Outcome		Impact		Summary															
Condoms																					
Number of condoms distributed and sold / year (in millions)	120	Condom use with non-regular partners (%)	Women 15-49	67	New HIV infections (ages 15+, in 1000s)	<table border="1"> <thead> <tr> <th>Scores</th> </tr> </thead> <tbody> <tr> <td>AGYW&MPs</td> </tr> <tr> <td>4</td> </tr> <tr> <td>SW</td> </tr> <tr> <td>7</td> </tr> <tr> <td>MSM</td> </tr> <tr> <td>id</td> </tr> <tr> <td>PWID</td> </tr> <tr> <td>id</td> </tr> <tr> <td>Condoms</td> </tr> <tr> <td>8</td> </tr> <tr> <td>VMMC</td> </tr> <tr> <td>6</td> </tr> <tr> <td>PrEP</td> </tr> <tr> <td>8</td> </tr> </tbody> </table>	Scores	AGYW&MPs	4	SW	7	MSM	id	PWID	id	Condoms	8	VMMC	6	PrEP	8
Scores																					
AGYW&MPs																					
4																					
SW																					
7																					
MSM																					
id																					
PWID																					
id																					
Condoms																					
8																					
VMMC																					
6																					
PrEP																					
8																					
Number of condoms distributed and sold per man 15-64 / year	26	Men 15-49	85																		
% of condom distribution need met	93																				
Voluntary medical male circumcision (VMMC)																					
Number of VMMCs performed / year (in thousands)	326	% of 2020 VMMC target achieved	Men	30																	
% of annual VMMC target achieved	63	National MC prevalence (%)	Men 15-24	19																	
			Men 15-49	14																	
ARV-based prevention																					
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id																	
Number of people who received PrEP at least once in the past 12 months	5000	% of PLHIV on ART	All PLHIV	88																	
		% of PLHIV virally suppressed	All PLHIV	62																	
Key populations																					
<i>Sex workers (SW)</i>			reported by		HIV prevalence																
Population size estimate for sex workers (in thousands)	45	Condom use / last paid sex (%)	Sex workers	96	Sex workers <25 years	21															
% of SWs who received at least two HIV prevention interventions (past 3 mo)	44	Condom use / last paid sex (%)	Men 15-49	90	Sex workers all ages	41															
<i>Men who have sex with men (MSM)</i>																					
Population size estimate for men who have sex with men (in thousands)	id	Condom use / last anal sex (%)	MSM	id	MSM <25 years	id															
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id				MSM all ages	31															
<i>People who inject drugs (PWID)</i>																					
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id															
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% of PWID on OST	PWID	id	PWID all ages	id															
<i>Structural barriers and enablers</i>																					
Training on HIV-related rights & non-discrimination: law enforcement officials	Limited	% of people who avoided health care because of stigma and discrimination	Sex workers	39																	
Training on HIV-related rights & non-discrimination: health care workers	National		MSM	id																	
			PWID	id																	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings																					
% of priority locations/districts with dedicated programs for AGYW & partners	17	Condom use with non-regular partners (%)	YW 15-24	57	Young women 15-24	5.7															
% of girls who completed lower secondary education	74		YM 15-24	84	Young men 15-24	3.2															
Sexual and reproductive health and rights strategy: Level of HIV linkages (%)	81				Adults 15-49	12.7															
<i>Background population-level factors</i>																					
% of women 15-49 experienced physical or sexual violence f. husband/partner	20	% who had multiple sexual partners	Men 15-49	14																	
% of married women who decide on their own health care	34																				
Acronyms																					
na ... not applicable		Very good			Low																
OST ... opioid substitution therapy		Good			Very low																
		Medium		id	Insufficient data																