

# Angola

2020

Based on most recent available data.

Version

2.01.

## Output (coverage)

## Outcome (service use/behaviour)

## Impact

## Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary														
<b>Condoms</b>																				
Number of condoms distributed and sold / year (in millions)	54	Condom use with non-regular partners (%)	Women 15-49	29	New HIV infections (ages 15+, in 1000s)	<table border="1"> <tr><td>AGYW&amp;MPs</td><td>3</td></tr> <tr><td>SW</td><td>5</td></tr> <tr><td>MSM</td><td>id</td></tr> <tr><td>PWID</td><td>id</td></tr> <tr><td>Condoms</td><td>3</td></tr> <tr><td>VMMC</td><td>na</td></tr> <tr><td>PrEP</td><td>0</td></tr> </table>	AGYW&MPs	3	SW	5	MSM	id	PWID	id	Condoms	3	VMMC	na	PrEP	0
AGYW&MPs	3																			
SW	5																			
MSM	id																			
PWID	id																			
Condoms	3																			
VMMC	na																			
PrEP	0																			
Number of condoms distributed and sold per man 15-64 /year	8	Men 15-49	53																	
% of condom distribution need met for men 15-64	25																			
<b>Voluntary medical male circumcision (VMMC)</b>																				
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na																
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	96																
			Men 15-49	96																
<b>ARV-based prevention</b>																				
Composite PrEP score (0-10)	0	% of national PrEP need met	All pop.	id	<table border="1"> <tr><td>HIV prevalence</td><td>na</td></tr> </table>	HIV prevalence	na													
HIV prevalence	na																			
Number of people who received PrEP at least once in the past 12 months	0	% of PLHIV virally suppressed	Women 15+	id																
% of PLHIV on ART	27	% of PLHIV virally suppressed	Men 15+	id																
<b>Key populations</b>																				
<i>Sex workers (SW)</i>																				
Population size estimate for sex workers (in thousands)	125	Condom use / last paid sex (%)	Sex workers	72	Sex workers <25 years	7														
% of SWs who received at least two HIV prevention interventions (past 3 mo)	43	Condom use / last paid sex (%)	Men 15-49	71	Sex workers all ages	8														
Prevention strategy includes core elements of SW prevention package	> Half	% of PLHIV on ART	SWs LHIV	42																
<i>Gay men and other men who have sex with men (MSM)</i>																				
Population size estimate for men who have sex with men (in thousands)	106	Condom use / last anal sex (%)	MSM	59	MSM <25 years	8														
% of MSM who received at least two HIV prevention interventions (past 3 mo)	id	% of PLHIV on ART	MSM LHIV	34	MSM all ages	2														
Prevention strategy includes core elements of MSM prevention package	> Half																			
<i>People who inject drugs (PWID)</i>																				
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id														
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id														
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id																
<i>Structural barriers and enablers</i>																				
Criminalization of selling sex	No	% of people who avoided health care because of stigma and discrimination	Sex workers	id																
Criminalization of same sex relations	No		MSM	id																
Criminalization of drug use/consumption or possession for personal use	id		PWID	id																
<b>Adolescent girls, young women (AGYW) &amp; partners in high-HIV incidence settings</b>																				
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	YW 15-24	33	Young women 15-24	1.2														
% of girls who completed lower secondary education	31		YM 15-24	52	Young men 15-24	0.4														
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	18	Adults 15-49	1.9														
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	24																
Provider-initiated condom promotion integrated into SRH services	Yes		Women 15-49	26																
HIV testing services integrated with SRH services	Yes																			

**Acronyms** na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV;YW...young women;YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data