

Cameroon

2020

Based on most recent available data.

Version

2.01.

Output (coverage)

Outcome (service use/behaviour)

Impact

Summary

Condoms

Number of condoms distributed and sold / year (in millions)	38
Number of condoms distributed and sold per man 15-64 /year	5
% of condom distribution need met for men 15-64	14

Condom use with non-regular partners (%)	Women 15-49	43
	Men 15-49	63

Voluntary medical male circumcision (VMMC)


Number of VMMCs performed / year (in thousands)	na
% of annual VMMC target achieved	na

% of 2020 VMMC target achieved	Men	na
National male circumcision prevalence (%)	Men 15-24	92
	Men 15-49	93

ARV-based prevention

Composite PrEP score (0-10)	1
Number of people who received PrEP at least once in the past 12 months	400
% of PLHIV on ART	62

% of national PrEP need met	All pop.	id
% of PLHIV virally suppressed	Women 15+	id
% of PLHIV virally suppressed	Men 15+	id



Key populations

Sex workers (SW)

Population size estimate for sex workers (in thousands)	71
% of SWs who received at least two HIV prevention interventions (past 3 mo)	23
Prevention strategy includes core elements of SW prevention package	All

Condom use / last paid sex (%)	Sex workers	91
Condom use / last paid sex (%)	Men 15-49	83
% of PLHIV on ART	SWs LHIV	99

Sex workers <25 years	id
Sex workers all ages	24

Gay men and other men who have sex with men (MSM)

Population size estimate for men who have sex with men (in thousands)	7
% of MSM who received at least two HIV prevention interventions (past 3 mo)	17
Prevention strategy includes core elements of MSM prevention package	All

Condom use / last anal sex (%)	MSM	77
% of PLHIV on ART	MSM LHIV	97

MSM <25 years	id
MSM all ages	21

People who inject drugs (PWID)

Population size estimate for people who inject drugs (in thousands)	2
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id
Prevention strategy includes core elements of PWID harm reduction package	None

% with safe injecting practices	PWID	id
% on opioid substitution therapy	PWID	id
% of PLHIV on ART	PWID LHIV	id

PWID <25 years	id
PWID all ages	id

Structural barriers and enablers

Criminalization of selling sex	Yes
Criminalization of same sex relations	Yes
Criminalization of drug use/consumption or possession for personal use	Yes

% of people who avoided health care because of stigma and discrimination	Sex workers	5
	MSM	14
	PWID	id

Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings

% of priority locations/districts with dedicated programs for AGYW & partners	5
% of girls who completed lower secondary education	68
Educational policies on HIV & sexuality education (secondary school)	Yes
Laws requiring parental consent for adolescents to access SRH services	No
Provider-initiated condom promotion integrated into SRH services	Yes
HIV testing services integrated with SRH services	Yes

Condom use with non-regular partners (%)	YW 15-24	50
	YM 15-24	65
% who had multiple sexual partners	Men 15-49	23
% of women experienced physical or sexual violence from husband/partner	YW 15-19	9
	Women 15-49	22

Young women 15-24	1.4
Young men 15-24	0.6
Adults 15-49	3.1

Scores

AGYW&MPs

3

SW

5

MSM

5

PWID

id

Condoms

3

VMMC

na

PrEP

1

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV;YW...young women;YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data