

Output (coverage)

Outcome (service use/behaviour)

Impact

Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary
Condoms				New HIV infections (ages 15+, in 1000s)		Scores
Number of condoms distributed and sold / year (in millions)	15	Condom use with non-regular partners (%)	Women 15-49	54		AGYW&MPs
Number of condoms distributed and sold per man 15-64 /year	38		Men 15-49	67		7
% of condom distribution need met for men 15-64	77					SW
Voluntary medical male circumcision (VMMC)						4
Number of VMMCs performed / year (in thousands)	17	% of 2020 VMMC target achieved	Men	35		MSM
% of annual VMMC target achieved	48	National male circumcision prevalence (%)	Men 15-24	12		3
			Men 15-49	10		PWID
						3
ARV-based prevention						Condoms
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id		7
Number of people who received PrEP at least once in the past 12 months	5607	% of PLHIV virally suppressed	Women 15+	97		VMMC
% of PLHIV on ART	96	% of PLHIV virally suppressed	Men 15+	87		5
Key populations				HIV prevalence		PrEP
<i>Sex workers (SW)</i>						8
Population size estimate for sex workers (in thousands)	15	Condom use / last paid sex (%)	Sex workers	83	Sex workers <25 years	64
% of SWs who received at least two HIV prevention interventions (past 3 mo)	14	Condom use / last paid sex (%)	Men 15-49	id	Sex workers all ages	61
Prevention strategy includes core elements of SW prevention package	All	% of PLHIV on ART	SWs LHIV	id		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	6	Condom use / last anal sex (%)	MSM	58	MSM <25 years	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	10	% of PLHIV on ART	MSM LHIV	id	MSM all ages	13
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	1	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	31	% on opioid substitution therapy	PWID	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id		
<i>Structural barriers and enablers</i>						
Criminalization of selling sex	Partial	% of people who avoided health care because of stigma and discrimination	Sex workers	21		
Criminalization of same sex relations	Yes		MSM	35		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	76	Condom use with non-regular partners (%)	YW 15-24	71	Young women 15-24	12.3
% of girls who completed lower secondary education	45		YM 15-24	93	Young men 15-24	4.1
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	14	Adults 15-49	27.0
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	id		
Provider-initiated condom promotion integrated into SRH services	No/id		Women 15-49	id		
HIV testing services integrated with SRH services	Yes					

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV;YW...young women;YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data