

HIV prevention among key populations - thematic

2020

Version 2.01

Thematic area	Indicator	Source	Angola	Botswana	Cameroon	Cote d'Ivoire	DR Congo	Ethiopia	Ghana	Kenya	Lesotho	Malawi	Mozambique	Namibia	Nigeria	South Africa	eswatini	Tanzania	Uganda	Zambia	Zimbabwe	Brazil	China	India	Indonesia	Iran	Mexico	Myanmar	Pakistan	Ukraine	
Impact	HIV prevalence (sex workers, <25 years)	GAM 3.3A/ IBBS	7	id	id	6	5	id	3	id	id	id	id	14	10	id	64	id	id	id	25	2	0	1	1	1	5	4	2	1	
	HIV prevalence (sex workers, all ages)	GAM 3.3A/ IBBS	8	42	24	13	8	24	5	29	72	55	id	28	14	58	61	15	31	49	42	5	0	2	2	2	1	8	2	6	
	HIV prevalence (men who have sex with men, <25 years)	GAM 3.3B/ IBBS	8	id	id	16	2	id	id	12	id	id	id	4	19	id	id	id	9	10	6	13	id	13	6	4	7				
	HIV prevalence (men who have sex with men, all ages)	GAM 3.3B/ IBBS	2	15	21	12	7	id	18	18	33	7	id	9	23	18	13	8	13	id	21	18	6	2	18	id	12	9	5	8	
	HIV prevalence (people who inject drugs, <25 years)	GAM 3.3C/ IBBS	id	id	id	0	1	id	id	id	id	id	id	id	2	12	id	12	17	id	id	na	3	8	1	0	4	17	18	5	
	HIV prevalence (people who inject drugs, all ages)	GAM 3.3C/ IBBS	id	id	id	6	4	id	id	18	id	id	id	id	3	22	id	16	17	id	id	na	6	6	14	4	3	19	38	24	
	HIV prevalence (transgender people, <25 years, %)	GAM 3.3D/ IBBS	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	18	19	id	6	19	0	id	id	6	id
HIV prevalence (transgender people, all ages, %)	GAM 3.3D/ IBBS	id	16	id	24	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	28	30	id	3	25	2	9	id	7	id	
Outcome	Condom use of sex workers with most recent client, % (reported by sex workers)	GAM 3.6A/ IBBS	72	76	91	76	74	98	90	92	62	65	id	86	98	86	83	72	69	79	75	86	93	91	67	59	87	90	51	94	
	Condom use at last anal sex among MSM (%)	GAM 3.6B/ IBBS	59	78	77	75	51	id	48	79	46	44	id	id	51	98	58	14	39	id	69	64	86	84	70	id	73	57	26	78	
	Condom use among transgender people (%)	GAM 3.6D/ IBBS	id	60	id	36	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	71	id	80	id	43	75	id	28	id	
	Condom use at last paid sex act, % (reported by men)	DHS or other survey	71	id	83	42	34	81	44	74	90	75	31	67	74	83	id	44	73	56	90	id	id	48	33	id	id	77	id	id	
	% of PLHIV on ART – sex workers	GAM3.5A	42	88	99	22	id	id	99	73	id	81	id	34	16	24	id	id	65	86	78	id	id	id	id	id	id	59	5	29	
	% of PLHIV on ART – men who have sex with men	GAM3.5B	34	74	97	10	id	id	4	63	id	86	id	15	14	28	id	id	66	id	94	id	91	id	id	id	id	44	1	46	
	% of PLHIV on ART – people who inject drugs	GAM3.5C	id	id	id	1	id	id	68	id	id	id	id	id	25	id	id	id	78	id	id	na	82	id	id	44	id	id	16	38	
	Safe injecting practices (people who inject drugs, %)	IBBS/ GAM 3.8	id	id	id	id	51	id	id	88	id	id	id	id	71	id	id	84	id	id	id	na	91	86	90	73	71	91	58	97	
	% of opioid users who receive opioid substitution therapy	GAM	id	id	id	id	id	id	id	22	id	id	id	id	id	id	id	id	24	id	id	na	id	22	11	13	id	21	id	5	
Output	Sex workers																														
	Population size estimate for female sex workers in 1000s	GAM 3.2A/ Pop. size study	125	18	71	50	350	211	60	207	8	37	93	8	874	125	15	155	174	134	45	547	id	868	278	90	240	69	230	87	
	% of all sex workers who received at least two HIV prevention interventions in the past three months	GAM 3.7A/ IBBS/prg. data	43	40	23	77	id	id	50	88	27	50	51	60	18	45	14	58	42	47	44	22	94	88	40	36	id	50	1	40	
	% of sex workers who avoided health care because of stigma and discrimination	GAM	id	id	5	23	id	id	7	id	8	id	id	id	id	id	21	id	id	id	39	12	id	id	id	id	id	id	4	id	id
	Prevention strategy includes core elements of SW prevention package	NCPI	> Half	> Half	All	All	All	> Half	All	All	> Half	All	> Half	> Half	All	All	All	> Half	All	> Half	All	> Half	All	All	id	> Half	> Half	All	> Half	> Half	
	Criminalization of selling sex	NCPI	No	Partial	Yes	Partial	Partial	Partial	Yes	Yes	Yes	Partial	No	Yes	Partial	Yes	Partial	Yes	Yes	Partial	Yes	No	Partial	Yes	id	Yes	Partial	Yes	Yes	Yes	
	Men who have sex with men																														
	Population size estimate for men who have sex with men in 1000s	GAM 3.2B/ Pop. size study/	106	10	7	36	195	id	55	51	6	48	41	7	313	347	6	50	44	68	16	2000	id	357	503	243	1200	252	830	179	
	% of all MSM who who received at least two HIV prevention interventions in the past three months	GAM 3.7B/ IBBS/prg. data	id	42	17	38	id	id	55	79	49	10	4	64	20	33	10	28	12	13	26	30	91	73	20	id	id	29	1	11	
	% of men who have sex with men who avoided health care because of stigma and discrimination	GAM	id	id	14	22	id	id	id	id	8	id	id	id	id	id	35	id	id	id	8	63	id	id	id	id	id	id	8	id	id
	Prevention strategy includes core elements of MSM prevention package	NCPI	> Half	< Half	All	All	> Half	id	All	All	> Half	All	> Half	> Half	> Half	All	> Half	> Half	> Half	> Half	< Half	All	All	All	id	id	> Half	All	< Half	< Half	
	Criminalization of same sex relations	NCPI	No	No	Yes	No	No	Yes	Yes	Yes	No	Yes	No	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	No	No	No	No	Yes	No	Yes	Yes	No
	People who inject drugs																														
	Population size estimate for people who inject drugs available	GAM 3.2C/ Pop. size study/	id	id	2	0	id	id	id	20	id	id	14	id	227	75	1	36	7	27	id	na	id	177	35	187	109	95	110	350	
	% of all people who inject drugs who received at least two HIV prevention interventions in the past three months	GAM 3.7C/ IBBS/prg. data	id	id	id	16	id	id	id	89	id	id	15	id	9	24	31	21	8	id	id	na	54	89	44	26	id	34	2	48	
	Number of needles and syringes per person who inject drugs	GAM 3.9/ Prg. records	id	id	id	id	1	id	id	137	id	id	id	id	id	id	id	id	15	id	id	id	na	246	349	3	43	6	369	46	53
	% of people who inject drugs who avoided health care because of stigma and discrimination	GAM	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	64	id	id	na	id	id	id	id	id	id	id	id
Prevention strategy includes core elements of PWID harm reduction package	NCPI	None	None	None	Some	Some	None	None	Some	None	None	Some	None	None	All	None	Some	None	None	None	na	All	All	None	All	Some	Some	Some	Some		
Criminalization of drug use/consumption or possession for personal use	NCPI	id	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	id	id	No	Partial	id	Yes		
Transgender people																															
Population size estimate for transgender people	GAM 3.2D/ Pop. size study/	3	0	id	1	id	id	id	4	id	0	id	id	id	id	id	id	id	id	id	id	id	id	70	39	id	119	id	52	id	
% of all TG people who who received at least two HIV prevention interventions in the past three months	GAM 3.7D/ IBBS/prg. data	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	59	id	63	id	id	id	id	4	id	
% of transgender people who avoided health care because of stigma and discrimination	GAM	id	id	id	23	id	id	id	id	id	id	id	id	id	id	id	id	id	id	id	11	id	id	id	id	id	id	id	id	id	

Legend

Very good