

Malawi

2020

Based on most recent available data.

Version

2.01.

Output (coverage)

Outcome (service use/behaviour)

Impact

Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary
Condoms				New HIV infections (ages 15+, in 1000s)		Scores
Number of condoms distributed and sold / year (in millions)	104	Condom use with non-regular partners (%)	Women 15-49	49		AGYW&MPs
Number of condoms distributed and sold per man 15-64 / year	20		Men 15-49	73		4
% of condom distribution need met for men 15-64	95					5
Voluntary medical male circumcision (VMMC)						MSM
Number of VMMCs performed / year (in thousands)	114	% of 2020 VMMC target achieved	Men	19		3
% of annual VMMC target achieved	19	National male circumcision prevalence (%)	Men 15-24	29		PWID
			Men 15-49	28		id
						Condoms
ARV-based prevention						8
Composite PrEP score (0-10)	4	% of national PrEP need met	All pop.	id		VMMC
Number of people who received PrEP at least once in the past 12 months	459	% of PLHIV virally suppressed	Women 15+	81		2
% of PLHIV on ART	79	% of PLHIV virally suppressed	Men 15+	62		PrEP
Key populations				HIV prevalence		4
<i>Sex workers (SW)</i>						
Population size estimate for sex workers (in thousands)	37	Condom use / last paid sex (%)	Sex workers	65	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	50	Condom use / last paid sex (%)	Men 15-49	75	Sex workers all ages	55
Prevention strategy includes core elements of SW prevention package	All	% of PLHIV on ART	SWs LHIV	81		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	48	Condom use / last anal sex (%)	MSM	44	MSM <25 years	5
% of MSM who received at least two HIV prevention interventions (past 3 mo)	10	% of PLHIV on ART	MSM LHIV	86	MSM all ages	7
Prevention strategy includes core elements of MSM prevention package	All					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id		
<i>Structural barriers and enablers</i>						
Criminalization of selling sex	Partial	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same sex relations	Yes		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	28	Condom use with non-regular partners (%)	YW 15-24	53	Young women 15-24	4.2
% of girls who completed lower secondary education	32		YM 15-24	73	Young men 15-24	2.1
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	13	Adults 15-49	8.9
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	28		
Provider-initiated condom promotion integrated into SRH services	Yes		Women 15-49	41		
HIV testing services integrated with SRH services	Yes					

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data