


THE STATE OF HIV PREVENTION IN NIGERIA


2020

The number of adults newly infected with HIV declined from 90000 in 2010 to 82000 in 2019, a 10% decline.


Number of new HIV infections (≥15 years)



Change in new HIV infections




HIV prevalence




Policy and structural barriers

Key populations



Adolescent girls and young women



HIV prevention adaptations during COVID-19

Adaptation	Safe continuation of outreach	Online counselling
Young women	Yes	Yes
Key populations	Yes	Yes
Adopted multi-month dispensing	Yes	Expanded alternative access
Condoms	Yes	Yes
PrEP	Yes	No
HIV treatment	Yes	Yes
Safe injecting equipment	No	
Opioid substitution therapy (take home dosages)		No

Note: 'Yes' refers to the adaptation having been introduced (not necessarily it being universally available).

Linkages between HIV and sexual and reproductive health services

HIV testing services integrated within sexual and reproductive health	Yes
Provider-initiated condom promotion in family planning services	Yes

HIV programme coverage and outcomes

ADOLESCENT GIRLS, YOUNG WOMEN & MALE PARTNERS

Condom use with a non-regular partner among young people 15-24 years old (%)

Young women: 38, Young men: 62, Target: 90%

Received two prevention interventions in past 3 months (%): 18

Score: 4

KEY POPULATIONS

Sex workers: Condom use at last paid sex (%): 98, Clients: 74, Target: 95%

Gay men and other men who have sex with men (MSM): Condom use at last anal sex (%): 51, Target: 90%

People who inject drugs (PWID): Use of harm reduction services (%): 71, Target: 95%; On opioid substitution therapy: 40%, Target: 40%

Received two prevention interventions in past 3 months (%): 20

Score: 4

CONDOM PROGRAMMING

Condom use with a non-regular partner, 15-49 years (%): Women: 36, Men: 65, Target: 90%

Estimated condom distribution need met (%): 60

Score: 6

VOLUNTARY MEDICAL MALE CIRCUMCISION

Uptake of voluntary medical male circumcision: 99, Target: 90%

% of annual voluntary medical male circumcision target achieved: 99%

Score: na

ANTIRETROVIRAL DRUG-BASED PREVENTION

Pre-exposure prophylaxis: Number of PrEP users: 360 (2018), 118 (2019), 11413 (2020); Annual change (2019-20): +9572%

Antiretroviral treatment: People living with HIV virally suppressed: 73%, Target: 73%

Antiretroviral treatment coverage: Overall: 65%, Sex workers: 16%, MSM: 14%, PWID: 25%

Score: 5

Scores (1-10) Very good Good Medium Low Very low id ... insufficient data na ... not applicable

Data sources: UNAIDS 2020 HIV estimates; Global AIDS Monitoring 2020; Global HIV Prevention Coalition progress survey 2020; and ICF - the DHS Program STATcompiler. Data sources key populations coverage: Global AIDS Monitoring 2020, Global Fund Proposals 2020 and PEPFAR COP20. Note: the 2020 HIV estimates presented are for 2019. Other data points may refer to various years when the surveys were conducted.

Implementation of Prevention 2020 Roadmap

Ten actions	2017	2020
1 - HIV prevention needs assessment	Not done	Done
2 - HIV prevention targets	Not done	Done
3 - HIV prevention strategy	Done	Done
4 - Legal and policy reform	Not done	Partly done
5a - Key population size estimates	Not done	Partly done
5b - Defined key population package	Not done	Partly done
5c - Adolescent girls and young women size estimates	Not done	Partly done
5d - Adolescent girls and young women package	Not done	Done
6 - Capacity development and technical assistance plan	Not done	Not done
7 - Social contracting	Not done	Not done
8 - HIV prevention financial gap analysis	Not done	Done
9 - Strengthen programme monitoring	Not done	Done
10 - Performance review	Not done	Done

Legend: Done (Green), Partly done (Yellow), Not done (Orange)