

Select country from dropdown menu

Angola

AGO

Based on most recent available data.

Version

2.01.

2021

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms				New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)		
Number of condoms distributed and sold / year (in millions)	id	Condom use with non-regular partners (%)	Women 15-49	29	<p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>	
Number of condoms distributed/sold per couple-year (age range 15-64)	id		Men 15-49	53		
Estimated condom distribution need met (%)	id					
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	96		
			Men 15-49	96		
ARV-based prevention						
Composite PrEP score (0-10)	id	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	id	% of PLHIV virally suppressed	Women 15+	id		
% of PLHIV on ART	33	% of PLHIV virally suppressed	Men 15+	id		
Key populations				HIV prevalence		
Sex workers (SW)						
Population size estimate for sex workers	54000	Condom use / last paid sex (%)	Sex workers	72	Sex workers <25 years	7.2
% of SWs who received at least two HIV prevention interventions (past 3 mo)	11	Condom use / last paid sex (%)	Men 15-49	71	Sex workers all ages	8.0
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	42		
Gay men and other men who have sex with men (MSM)						
Population size estimate for men who have sex with men	29400	Condom use / last anal sex (%)	MSM	59	MSM <25 years	8.2
% of MSM who received at least two HIV prevention interventions (past 3 mo)	3	% on ART	MSM LHIV	id	MSM all ages	2.0
Prevention strategy includes core elements of MSM prevention package	> Half					
People who inject drugs (PWID)						
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id		
Structural barriers and enablers						
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same-sex sexual acts	No		MSM	id		
Criminalization of drug use/consumption or possession for personal use	id		PWID	id		
Criminalization of transgender people	Yes		Transgender	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	YW 15-24	33	Young women 15-24	1.0
Educational policies on HIV & sexuality education (secondary school)	Yes		YM 15-24	52	Young men 15-24	0.4
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	32	Adults 15-49	1.7
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	24		
HIV testing services integrated with SRH services	Yes		Women 15-49	26		

2010 baseline
2020
2020 target
2025 target

2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

Summary Scores
AGYW&MPs
3
SW
3
MSM
3
PWID
id
Condoms
id
VMMC
na
ART
3
PrEP
id

Acronyms na ... not applicable;
PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
SRH ... sexual and reproductive health.

Very good	Low
Good	Very low
Medium	id
	Insufficient data