

Select country from dropdown menu

Botswana

BWA

Based on most recent available data.

Version

2.01.

2021

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)			Outcome (service use/behaviour)			Impact			
Condoms						<p>New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)</p> <p>2010 baseline: 13 000 2020: 8 600 2020 target: 3 200 2025 target: 2 500</p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>			
Number of condoms distributed and sold / year (in millions)	27		Condom use with non-regular partners (%)	Women 15-49	id				
Number of condoms distributed/sold per couple-year (age range 15-64)	38			Men 15-49	id				
Estimated condom distribution need met (%)	id								
Voluntary medical male circumcision (VMMC)									
Number of VMMCs performed / year (in thousands)	3		% of 2020 VMMC target achieved	Men	29				
% of annual VMMC target achieved	6		National male circumcision prevalence (%)	Men 15-24	23				
				Men 15-49	26				
ARV-based prevention									
Composite PrEP score (0-10)	5		% of national PrEP need met	All pop.	id				
Number of people who received PrEP at least once in the past 12 months	2259		% of PLHIV virally suppressed	Women 15+	93				
% of PLHIV on ART	87		% of PLHIV virally suppressed	Men 15+	76				
Key populations						HIV prevalence			
<i>Sex workers (SW)</i>									
Population size estimate for sex workers	6700		Condom use / last paid sex (%)	Sex workers	76	Sex workers <25 years	id	Summary Scores	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	37		Condom use / last paid sex (%)	Men 15-49	id	Sex workers all ages	42.2		
Prevention strategy includes core elements of SW prevention package	> Half		% on ART	SWs LHIV	88				
<i>Gay men and other men who have sex with men (MSM)</i>									
Population size estimate for men who have sex with men	2600		Condom use / last anal sex (%)	MSM	78	MSM <25 years	id		
% of MSM who received at least two HIV prevention interventions (past 3 mo)	50		% on ART	MSM LHIV	74	MSM all ages	14.8		
Prevention strategy includes core elements of MSM prevention package	< Half								
<i>People who inject drugs (PWID)</i>									
Population size estimate for people who inject drugs	id		% with safe injecting practices	PWID	id	PWID <25 years	id		
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id		% on opioid substitution therapy	PWID	id	PWID all ages	id		
Prevention strategy includes core elements of PWID harm reduction package	None		% on ART	PWID LHIV	id				
<i>Structural barriers and enablers</i>									
Criminalization of sex work	Yes		% of people who avoided health care because of stigma and discrimination	Sex workers	id				
Criminalization of same-sex sexual acts	No			MSM	id				
Criminalization of drug use/consumption or possession for personal use	Yes			PWID	id				
Criminalization of transgender people	No			Transgender	id				
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings									
% of priority locations/districts with dedicated programs for AGYW & partners	33		Condom use with non-regular partners (%)	YW 15-24	id	Young women 15-24	8.8		
Educational policies on HIV & sexuality education (secondary school)	Yes			YM 15-24	id	Young men 15-24	4.2		
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16		% who completed lower secondary education	Girls	92	Adults 15-49	21.7		
Provider-initiated condom promotion integrated into SRH services	Yes		% of women experienced physical or sexual violence from husband/partner	YW 15-19	id				
HIV testing services integrated with SRH services	Partial			Women 15-49	id				

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good
Good
Medium

Low
Very low
Insufficient data