

Select country from dropdown menu

Eswatini

SWZ

Based on most recent available data.

Version 2.01.

2021

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)			Outcome (service use/behaviour)			Impact		
Condoms						New HIV infections (ages 15+ trend vs. 2020 and 2025 targets) 		
Number of condoms distributed and sold / year (in millions)	15	Condom use with non-regular partners (%)	Women 15-49	54				
Number of condoms distributed/sold per couple-year (age range 15-64)	43		Men 15-49	67				
Estimated condom distribution need met (%)	86							
Voluntary medical male circumcision (VMMC)						HIV prevalence Sex workers <25 years 31.7 Sex workers all ages 60.8 MSM <25 years 7.3 MSM all ages 27.2 PWID <25 years id PWID all ages id		
Number of VMMCs performed / year (in thousands)	9	% of 2020 VMMC target achieved	Men	39				
% of annual VMMC target achieved	24	National male circumcision prevalence (%)	Men 15-24	5				
			Men 15-49	29				
ARV-based prevention						Summary Scores AGYW&MPs 7 SW 2 MSM 9 PWID 6 Condoms 7 VMMC 3 ART 10 PrEP 8		
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id				
Number of people who received PrEP at least once in the past 12 months	9125	% of PLHIV virally suppressed	Women 15+	>98				
% of PLHIV on ART	>98	% of PLHIV virally suppressed	Men 15+	89				
Key populations								
<i>Sex workers (SW)</i>								
Population size estimate for sex workers	4000	Condom use / last paid sex (%)	Sex workers	50				
% of SWs who received at least two HIV prevention interventions (past 3 mo)	30	Condom use / last paid sex (%)	Men 15-49	id				
Prevention strategy includes core elements of SW prevention package	< Half	% on ART	SWs LHIV	id				
<i>Gay men and other men who have sex with men (MSM)</i>								
Population size estimate for men who have sex with men	2400	Condom use / last anal sex (%)	MSM	80				
% of MSM who received at least two HIV prevention interventions (past 3 mo)	97	% on ART	MSM LHIV	id				
Prevention strategy includes core elements of MSM prevention package	< Half							
<i>People who inject drugs (PWID)</i>								
Population size estimate for people who inject drugs	300	% with safe injecting practices	PWID	id				
% of PWID who received at least two HIV prevention interventions (past 3 mo)	62	% on opioid substitution therapy	PWID	id				
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	PWID LHIV	id				
<i>Structural barriers and enablers</i>								
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id				
Criminalization of same-sex sexual acts	Yes		MSM	id				
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id				
Criminalization of transgender people	No		Transgender	id				
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								
% of priority locations/districts with dedicated programs for AGYW & partners	76	Condom use with non-regular partners (%)	YW 15-24	54		Young women 15-24	12.2	
Educational policies on HIV & sexuality education (secondary school)	Yes		YM 15-24	70		Young men 15-24	4.7	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	54		Adults 15-49	27.1	
Provider-initiated condom promotion integrated into SRH services	No/id	% of women experienced physical or sexual violence from husband/partner	YW 15-19	id				
HIV testing services integrated with SRH services	Yes		Women 15-49	id				

Acronyms na ... not applicable;
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
 SRH ... sexual and reproductive health.

 Very good	 Low
 Good	 Very low
 Medium	 Insufficient data