Select country from dropdown menu									
Mozambique	MOZ	Based on most recent available data.	Version	2.01.				2021	
Output (coverage)		Outcome (service use/behaviour)		\rightarrow		Impact			
Condoms Number of condoms distributed and sold / year (in millions) Number of condoms distributed/sold per couple-year (age range 15-64)	84 10	Condom use with non-regular partners (%)	Women 15-49 Men 15-49	42 47	140,000	New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)			2010 baseline 2020 2020 target 2025 target 2020 and 2025
Estimated condom distribution need met (%) Voluntary medical male circumcision (VMMC) Number of VMMCs performed / year (in thousands) % of annual VMMC target achieved	113 20	% of 2020 VMMC target achieved National male circumcision prevalence (%)	Men Men 15-24 Men 15-49	44 66 47	120,000 100,000 80,000 60,000 40,000		85 000	23 000	targets representhe country's required contribution to global targets, a 75% reduction b 2020 and 82.5%
ARV-based prevention Composite PrEP score (0-10)	4	% of national PrEP need met	All pop.	id	20,000		4 29 000	•	reduction by 2025 against
Number of people who received PrEP at least once in the past 12 months % of PLHIV on ART	18513 68	% of PLHIV virally surpressed % of PLHIV virally surpressed	Women 15+ Men 15+	60 50	2010	2015	2020	2025	2010 as a baseline.
Key populations	00	70 OFF EFFIC VITAITY SUI PICESSEE	Wich 151	30			HIV pi	revalence	Summary
Sex workers (SW)									Scores
Population size estimate for sex workers	224000	Condom use / last paid sex (%)	Sex workers	id		Sex workers <2	5 years	id	AGYW&M
% of SWs who received at least two HIV prevention interventions (past 3 mo)	19	Condom use / last paid sex (%)	Men 15-49	31		Sex workers	all ages	id	4
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id					sw
Gay men and other men who have sex with men (MSM)									3
Population size estimate for men who have sex with men	64000	Condom use / last anal sex (%)	MSM	id		MSM <2	5 years	id	MSM
% of MSM who received at least two HIV prevention interventions (past 3 mo)	21	% on ART	MSM LHIV	id		MSM a	all ages	id	id
Prevention strategy includes core elements of MSM prevention package	> Half								PWID
People who inject drugs (PWID)									0
Population size estimate for people who inject drugs	14000	% with safe injecting practices	PWID	id		PWID <2	5 years	id	Condom
% of PWID who received at least two HIV prevention interventions (past 3 mo)	3	% on opioid substitution therapy	PWID	id		PWID a	all ages	id	4
Prevention strategy includes core elements of PWID harm reduction package	Some	% on ART	PWID LHIV	id					VMMC
Structural barriers and enablers									3
Criminalization of sex work	No		Sex workers	id					ART
Criminalization of same-sex sexual acts	No	% of people who avoided health care because of	MSM	id					7
Criminalization of drug use/consumption or possession for personal use	Yes	stigma and discrimination	PWID	id					PrEP
Criminalization of transgender people	No		Transgender	id					4
dolescent girls, young women (AGYW) & partners in high-HIV incidence settings									
% of priority locations/districts with dedicated programs for AGYW & partners	42	Condom use with non-regular partners (%)	YW 15-24	51		Young womer		6.2	
Educational policies on HIV & sexuality education (secondary school)	Yes		YM 15-24	48		Young mer		2.4	
Laws requiring parental consent for adolescents to access HIV testing services		% who completed lower secondary education	Girls	11		Adults	5 15-49	11.3	
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual	YW 15-19	10					
HIV testing services integrated with SRH services	Yes	violence from husband/partner	Women 15-49	16					
cronyms na not applicable; LHIVpeople living with HIV; LHIVliving with HIV;YWyoung women;YMyoung men; RH sexual and reproductive health.		Very good Good Medium		id	Low Very low Insufficient dat	a			