

Select country from dropdown menu

Zimbabwe

ZWE

Based on most recent available data.

Version

2.01.

2021

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)			Outcome (service use/behaviour)			Impact		
Condoms						<p>New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)</p> <p>2010 baseline: 56 000 2020: 20 000 2020 target: 14 000 2025 target: 11 000</p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>		
Number of condoms distributed and sold / year (in millions)	83		Condom use with non-regular partners (%)	Women 15-49	65			
Number of condoms distributed/sold per couple-year (age range 15-64)	21			Men 15-49	82			
Estimated condom distribution need met (%)	67							
Voluntary medical male circumcision (VMMC)								
Number of VMMCs performed / year (in thousands)	82		% of 2020 VMMC target achieved	Men	45			
% of annual VMMC target achieved	16		National male circumcision prevalence (%)	Men 15-24	19			
				Men 15-49	14			
ARV-based prevention								
Composite PrEP score (0-10)	8		% of national PrEP need met	All pop.	id			
Number of people who received PrEP at least once in the past 12 months	48583		% of PLHIV virally suppressed	Women 15+	86			
% of PLHIV on ART	93		% of PLHIV virally suppressed	Men 15+	81			
Key populations						HIV prevalence		
<i>Sex workers (SW)</i>						Summary Scores		
Population size estimate for sex workers	45000		Condom use / last paid sex (%)	Sex workers	95	Sex workers <25 years	25.0	AGYW&MPs
% of SWs who received at least two HIV prevention interventions (past 3 mo)	35		Condom use / last paid sex (%)	Men 15-49	90	Sex workers all ages	42.2	4
Prevention strategy includes core elements of SW prevention package	All		% on ART	SWs LHIV	59			SW
<i>Gay men and other men who have sex with men (MSM)</i>								6
Population size estimate for men who have sex with men	23000		Condom use / last anal sex (%)	MSM	69	MSM <25 years	8.5	MSM
% of MSM who received at least two HIV prevention interventions (past 3 mo)	26		% on ART	MSM LHIV	73	MSM all ages	21.1	5
Prevention strategy includes core elements of MSM prevention package	All							PWID
<i>People who inject drugs (PWID)</i>								id
Population size estimate for people who inject drugs	id		% with safe injecting practices	PWID	id	PWID <25 years	id	Condoms
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id		% on opioid substitution therapy	PWID	id	PWID all ages	id	7
Prevention strategy includes core elements of PWID harm reduction package	None		% on ART	PWID LHIV	id			VMMC
<i>Structural barriers and enablers</i>								3
Criminalization of sex work	Yes			Sex workers	39			ART
Criminalization of same-sex sexual acts	Yes		% of people who avoided health care because of stigma and discrimination	MSM	8			9
Criminalization of drug use/consumption or possession for personal use	Yes			PWID	id			PrEP
Criminalization of transgender people	No			Transgender	11			8
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								
% of priority locations/districts with dedicated programs for AGYW & partners	18		Condom use with non-regular partners (%)	YW 15-24	54	Young women 15-24	5.2	
Educational policies on HIV & sexuality education (secondary school)	Yes			YM 15-24	81	Young men 15-24	2.9	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16		% who completed lower secondary education	Girls	53	Adults 15-49	12.6	
Provider-initiated condom promotion integrated into SRH services	Yes		% of women experienced physical or sexual violence from husband/partner	YW 15-19	31			
HIV testing services integrated with SRH services	Yes			Women 15-49	19			

Acronyms na ... not applicable;
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
 SRH ... sexual and reproductive health.

 Very good	 Low
 Good	 Very low
 Medium	 Insufficient data