

Select country from dropdown menu

Botswana

BWA

Based on most recent available data.

Version 2.02.

2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)			Outcome (service use/behaviour)			Impact		
Condoms						New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)		
Number of condoms distributed and sold / year (in millions)	17		Condom use with non-regular partners (%)	Women 15-49	id			
Number of condoms distributed/sold per couple-year (age range 15-64)	24			Men 15-49	id			
Estimated condom distribution need met (%)	id							
Voluntary medical male circumcision (VMMC)								
Number of VMMCs performed / year (in thousands)	4		% of 2020 VMMC target achieved	Men	29			
% of annual VMMC target achieved	6		National male circumcision prevalence (%)	Men 15-24	23			
				Men 15-49	26			
ARV-based prevention								
Composite PrEP score (0-10)	8		% of national PrEP need met	All pop.	id			
Number of people who received PrEP at least once in the past 12 months	9064		% of PLHIV virally suppressed	Women 15+	94			
% of PLHIV on ART	92			Men 15+	86			
Key populations						HIV prevalence		
<i>Sex workers (SW)</i>								
Population size estimate for sex workers	6700		Condom use / last paid sex (%)	Sex workers	76	Sex workers <25 years	id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	90			Men 15-49	id	Sex workers all ages	42.2	
Prevention strategy includes core elements of SW prevention package	> Half		% on ART	SWs LHIV	88			
<i>Gay men and other men who have sex with men (MSM)</i>								
Population size estimate for men who have sex with men	2600		Condom use / last anal sex (%)	MSM	78	MSM <25 years	id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	75		% on ART	MSM LHIV	74	MSM all ages	14.8	
Prevention strategy includes core elements of MSM prevention package	< Half							
<i>People who inject drugs (PWID)</i>								
Population size estimate for people who inject drugs	id		% with safe injecting practices	PWID	id	PWID <25 years	id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id		% on opioid substitution therapy	PWID	id	PWID all ages	id	
Prevention strategy includes core elements of PWID harm reduction package	None		% on ART	PWID LHIV	id			
<i>Structural barriers and enablers</i>								
Criminalization of sex work	Yes		% of people who avoided health care because of stigma and discrimination	Sex workers	id			
Criminalization of same-sex sexual acts	No			MSM	id			
Criminalization of drug use/consumption or possession for personal use	Yes			PWID	id			
Criminalization of transgender people	No			Transgender	id			
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings								
% of priority locations/districts with dedicated programs for AGYW & partners	56		Condom use with non-regular partners (%)	Women 15-24	id	Women 15-24	7.1	
Educational policies on HIV & sexuality education (secondary school)	Yes			Men 15-24	id	Men 15-24	3.8	
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <16		% who completed lower secondary education	Girls	92	Adults, 15-49	21.2	
Provider-initiated condom promotion integrated into SRH services	Yes		% of women experienced physical or sexual violence from husband/partner	Women 15-19	id			
HIV testing services integrated with SRH services	Partial			Women 15-49	id			

Summary Scores

AGYW	7
Sex workers	7
MSM	8
PWID	id
Condoms	id
VMMC	2
ART	9
PrEP	8

Acronyms: na ... not applicable;
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;
 SRH ... sexual and reproductive health.

 Very good	 Low
 Good	 Very low
 Medium	 Insufficient data